



JCO07 Rec'd PCT/PTO 12 MAR 2002

10/070760

TRANSMITTAL LETTER TO THE UNITED STATES PATENT & TRADEMARK OFFICE ATTORNEY'S DOCKET NUMBER 50730

DESIGNATED/ELECTED OFFICE (DO/EO/US)  
CONCERNING A FILING UNDER 35 U.S.C. 371

U.S. APPLICATION NO. (If known, see 37 CFR 1.5)

INTERNATIONAL APPLICATION NO.	INTERNATIONAL FILING DATE	PRIORITY DATE CLAIMED
PCT/EP00/09000	14 September 2000	15 September 1999

TITLE OF INVENTION: UNSATURATED OXIME ETHERS AND THE USE THEREOF FOR CONTROL OF HARMFUL FUNGI AND VETERINARY PESTS

APPLICANT(S) FOR DO/EO/US Markus GEWEHR, Hubert SAUTER, Bernd MUELLER, Andreas GYPSEER, Wassilios GRAMMENOS, Jordi Tormo i BLASCO, Arne PTOCK, Oliver CULLMANN, Thomas GROTE, Eberhard AMMERMAN, Siegfried STRATHMANN, Gisela LORENZ, Volker HARRIES

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. /X/ This is a FIRST submission of items concerning a filing under 35 U.S.C. 371.
2. / / This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371.
3. /X/ This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).
4. /X/ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
5. /X/ A copy of the International Application as filed (35 U.S.C. 371(c)(2)).
  - a./X/ is transmitted herewith (required only if not transmitted by the International Bureau).
  - b./ / has been transmitted by the International Bureau.
  - c./ / is not required, as the application was filed in the United States Receiving Office (RO/USO).
6. /X/ A translation of the International Application into English (35 U.S.C. 371(c)(2)).
7. /X / Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3)).
  - a./X / are transmitted herewith (required only if not transmitted by the International Bureau).
  - b./ / have been transmitted by the International Bureau.
  - c./ / have not been made; however, the time limit for making such amendments has NOT expired.
  - d./ / have not been made and will not be made.
8. /X / A translation of the amendments to the claims under PCT Article 19(35 U.S.C. 371(c)(3)).
9. /X / An oath or declaration of the inventor(s)(35 U.S.C. 371(c)(4)).
- 10./ / A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11. to 16. below concern other document(s) or information included:

- 11./ / An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
- 12./X / An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
- 13./X / A FIRST preliminary amendment.  
/ / A SECOND or SUBSEQUENT preliminary amendment.
- 14./ / A substitute specification.
- 15./ / A change of power of attorney and/or address letter.
- 16./X / Other items or information.  
International Search Report  
International Preliminary Examination Report

10/070760

JC10 Rec'd PCT/PTO 12 MAR 2002

U.S. Appl. No. (If Known) INTERNATIONAL APPL. NO.  
PCT/EP00/0900ATTORNEY'S DOCKET NO.  
50730

		CALCULATIONS	PTO USE ONLY
17. /X/ The following fees are submitted			
BASIC NATIONAL FEE (37 CFR 1.492(a)(1)-(5)):			
Search Report has been prepared by the			
EPO or JPO.....	\$890.00	890.00	
International preliminary examination fee paid to USPTO			
(37 CFR 1.482).....	\$710.00		
No international preliminary examination fee paid to			
USPTO (37 CFR 1.482) but international search fee paid			
to USPTO (37 CFR 1.445(a)(2)).....\$740.00			
Neither international preliminary examination fee			
(37 CFR 1.482) nor international search fee			
(37 CFR 1.445(a)(2)) paid to USPTO .....\$ 1,040.00			
International preliminary examination fee paid to			
USPTO (37 CFR 1.482) and all claims satisfied pro			
-visions of PCT Article 33(2)-(4).....\$100.00			
ENTER APPROPRIATE BASIC FEE AMOUNT =		\$ 890.00	
Surcharge of \$130.00 for furnishing the oath or declaration			
later than // 20 // 30 months from the earliest			
claimed priority date (37 CFR 1.492(e)).			
Claims	Number Filed	Number Extra	Rate
Total Claims	11 -20		X\$18.
Indep. Claims	2 -3		X\$84.
Multiple dependent claim(s) (if applicable)			+280.
TOTAL OF ABOVE CALCULATION		=	890.
Reduction of 1/2 for filing by small entity, if applicable.			
Verified Small Entity statement must also be filed			
(Note 37 CFR 1.9, 1.27, 1.28).			
SUBTOTAL		=	890.
Processing fee of \$130. for furnishing the English			
translation later than // 20 // 30 months from the			
earliest claimed priority date (37 CFR 1.492(f)).			
TOTAL NATIONAL FEE		=	890.
Fee for recording the enclosed assignment (37 CFR 1.21(h)).			
The assignment must be accompanied by an appropriate cover			
sheet (37 CFR 3.28, 3.31) \$40.00 per property =			
TOTAL FEES ENCLOSED		=	\$ 930.00
		Amount to be	
		refunded:	\$
		Charged	\$

a./X/ A check in the amount of \$ 930.00 to cover the above fees is enclosed.

b./ / Please charge my Deposit Account No. \_\_\_\_\_ in the amount of \$ \_\_\_\_\_ to cover the above fees. A duplicate copy of this sheet is enclosed.

c./X/ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 11-0345. A duplicate copy of this sheet is enclosed.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:

KEIL & WEINKAUF  
1101 Connecticut Ave., N.W.  
Washington, D. C. 20036

SIGNATURE

Herbert B. Keil  
NAME

Registration No. 18,967

10/070760

JC10 Rec'd PCT/PTO 12 MAR 2002

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re the Application of )  
GEWEHR et al. ) BOX PCT  
)  
International Application )  
PCT/EP 00/00900 )  
)  
Filed: September 14, 2000 )  
)

For: UNSATURATED OXIME ETHERS AND THE USE THEREOF FOR CONTROL OF  
HARMFUL FUNGI AND BETERINARY PESTS

PRELIMINARY AMENDMENT

Honorable Commissioner of  
Patents and Trademarks  
Washington, D.C. 20231

Sir:

Prior to examination, kindly amend the above-identified application as follows:

IN THE CLAIMS

Kindly amend the claims as shown on the attached sheets.

R E M A R K S

The claims have been amended to eliminate multiple dependency and to place them in  
better form for U.S. filing. No new matter is included.

A clean copy of the claims is attached.

Favorable action is solicited.

Respectfully submitted,

KEIL & WEINKAUF



Herbert B. Keil  
Reg. No. 18,967

1101 Connecticut Ave., N.W.  
Washington, D.C. 20036

(202)659-0100

CLEAN VERSION OF AMENDED CLAIMS - OZ 50730

3. A compound of the formula 1 as claimed in claim 1 where:

W is  $-\text{OCH}_2-$ ,  $-\text{C}(\text{R}_{10})=\text{N}-\text{O}-\text{CH}_2$ ;

X is halogen,  $\text{C}_1$ - $\text{C}_4$ -alkoxy;

$\text{R}_1$  is H,  $\text{C}_1$ - $\text{C}_4$ -alkyl;

$\text{R}_2$  is H,  $\text{C}_1$ - $\text{C}_4$ -alkyl;

n is 1 or 2;

$\text{R}_3$  is  $\text{C}_1$ - $\text{C}_4$ -alkyl;

$\text{R}_5$  is H,  $\text{C}_1$ - $\text{C}_4$ -alkyl;

$\text{R}_6$  is H,  $\text{C}_1$ - $\text{C}_4$ -alkyl;

$\text{R}_7$  is H, halogen,  $\text{C}_1$ - $\text{C}_6$ -alkyl;

$\text{R}_8$  is H, halogen,  $\text{C}_1$ - $\text{C}_6$ -alkyl,  $\text{C}_2$ - $\text{C}_6$ -alkenyl; or

$\text{R}_7$  and  $\text{R}_8$ , together with the carbon atoms to which they are attached, form a thiophenyl, furanyl, oxazolyl or thiazolyl radical, where these groups may have one or two substituents which are selected independently of one another from the group consisting of  $\text{C}_1$ - $\text{C}_4$ -alkyl, halogen and phenyl which may be substituted by one or two halogen;

$\text{R}_9$  is H, halogen,  $\text{C}_1$ - $\text{C}_6$ -alkyl;

$\text{R}_{10}$  is H,  $\text{C}_1$ - $\text{C}_4$ -alkyl.

4. A compound of the formula 1 as claimed in claim 1 where:

W is  $-\text{OCH}_2-$ ,  $-\text{C}(\text{R}_{10})=\text{N}-\text{O}-\text{CH}_2$ ;

X is  $\text{C}_1$ - $\text{C}_4$ -alkoxy;

CLEAN VERSION OF AMENDED CLAIMS - 50730

R<sub>1</sub> is H;

R<sub>2</sub> is H, C<sub>1</sub>-C<sub>4</sub>-alkyl;

n is 1 or 2;

R<sub>3</sub> is C<sub>1</sub>-C<sub>4</sub>-alkyl;

R<sub>5</sub> is H, C<sub>1</sub>-C<sub>4</sub>-alkyl;

R<sub>6</sub> is H, C<sub>1</sub>-C<sub>4</sub>-alkyl;

R<sub>7</sub> is H, halogen;

R<sub>8</sub> is H, C<sub>1</sub>-C<sub>4</sub>-alkyl, halogen; or

R<sub>7</sub> and R<sub>8</sub>, together with the carbon atoms to which they are attached, form a thiophenyl or oxazolyl radical, where these groups may be substituted by one or two halogen or phenyl and the phenyl may be substituted by one or two halogen;

R<sub>9</sub> is H, halogen;

R<sub>10</sub> is H, C<sub>1</sub>-C<sub>4</sub>-alkyl.

6. The use of the compounds of the formula 1 as claimed in claim 1 as fungicides or for controlling pests.

3. A compound of the formula 1 as claimed in claim 1 [or 2]  
where:

W is  $-\text{OCH}_2-$ ,  $-\text{C}(\text{R}_{10})=\text{N}-\text{O}-\text{CH}_2$ ;

X is halogen,  $\text{C}_1$ - $\text{C}_4$ -alkoxy;

$\text{R}_1$  is H,  $\text{C}_1$ - $\text{C}_4$ -alkyl;

$\text{R}_2$  is H,  $\text{C}_1$ - $\text{C}_4$ -alkyl;

n is 1 or 2;

$\text{R}_3$  is  $\text{C}_1$ - $\text{C}_4$ -alkyl;

$\text{R}_5$  is H,  $\text{C}_1$ - $\text{C}_4$ -alkyl;

$\text{R}_6$  is H,  $\text{C}_1$ - $\text{C}_4$ -alkyl;

$\text{R}_7$  is H, halogen,  $\text{C}_1$ - $\text{C}_6$ -alkyl;

$\text{R}_8$  is H, halogen,  $\text{C}_1$ - $\text{C}_6$ -alkyl,  $\text{C}_2$ - $\text{C}_6$ -alkenyl; or

$\text{R}_7$  and  $\text{R}_8$ , together with the carbon atoms to which they are attached, form a thiophenyl, furanyl, oxazolyl or thiazolyl radical, where these groups may have one or two substituents which are selected independently of one another from the group consisting of  $\text{C}_1$ - $\text{C}_4$ -alkyl, halogen and phenyl which may be substituted by one or two halogen;

$\text{R}_9$  is H, halogen,  $\text{C}_1$ - $\text{C}_6$ -alkyl;

$\text{R}_{10}$  is H,  $\text{C}_1$ - $\text{C}_4$ -alkyl.

4. A compound of the formula 1 as claimed in claim 1 [any of the preceding claims] where:

W is  $-\text{OCH}_2-$ ,  $-\text{C}(\text{R}_{10})=\text{N}-\text{O}-\text{CH}_2$ ;

X is  $\text{C}_1$ - $\text{C}_4$ -alkoxy;

MARKED VERSION OF AMENDED CLAIMS - 50730

R<sub>1</sub> is H;

R<sub>2</sub> is H, C<sub>1</sub>-C<sub>4</sub>-alkyl;

n is 1 or 2;

R<sub>3</sub> is C<sub>1</sub>-C<sub>4</sub>-alkyl;

R<sub>5</sub> is H, C<sub>1</sub>-C<sub>4</sub>-alkyl;

R<sub>6</sub> is H, C<sub>1</sub>-C<sub>4</sub>-alkyl;

R<sub>7</sub> is H, halogen;

R<sub>8</sub> is H, C<sub>1</sub>-C<sub>4</sub>-alkyl, halogen; or


R<sub>7</sub> and R<sub>8</sub>, together with the carbon atoms to which they are attached, form a thiophenyl or oxazolyl radical, where these groups may be substituted by one or two halogen or phenyl and the phenyl may be substituted by one or two halogen;

R<sub>9</sub> is H, halogen;

R<sub>10</sub> is H, C<sub>1</sub>-C<sub>4</sub>-alkyl.

6. The use of the compounds of the formula 1 as claimed in claim 1 [any of claims 1 to 5] as fungicides or for controlling pests.

1. An oxime ether compound of the formula 1



where:

W is  $-OCH_2-$ ,  $-C(R_{10})=N-O-CH_2-$ ;

X is halogen,  $C_1$ - $C_4$ -alkyl,  $C_1$ - $C_4$ -alkoxy;

$R_1$  is H,  $C_1$ - $C_4$ -alkyl, halogen, nitro, CN, halo- $C_1$ - $C_4$ -alkyl,  $C_1$ - $C_4$ -alkoxy;

$R_2$  is H,  $C_1$ - $C_4$ -alkyl, halogen, nitro, CN, halo- $C_1$ - $C_4$ -alkyl,  $C_1$ - $C_4$ -alkoxy;

n is 1 or 2;

$R_3$  is H,  $C_1$ - $C_4$ -alkyl;

$R_5$  is H,  $C_1$ - $C_4$ -alkyl,  $C_2$ - $C_4$ -alkenyl;

$R_6$  is H,  $C_1$ - $C_4$ -alkyl,  $C_1$ - $C_4$ -haloalkyl,  $C_2$ - $C_4$ -alkenyl, aryl;

$R_7$  is H, halogen,  $C_1$ - $C_6$ -alkyl,  $C_1$ - $C_6$ -haloalkyl,  $C_2$ - $C_6$ -alkenyl,  $C_2$ - $C_6$ -haloalkenyl,  $C_3$ - $C_6$ -cycloalkyl,  $C_3$ - $C_6$ -halocycloalkyl, unsubstituted or substituted aryl;

$R_8$  is H, halogen,  $C_1$ - $C_6$ -alkyl,  $C_1$ - $C_6$ -haloalkyl,  $C_2$ - $C_6$ -alkenyl,  $C_2$ - $C_6$ -haloalkenyl,  $C_3$ - $C_6$ -cycloalkyl,  $C_3$ - $C_6$ -halocycloalkyl, unsubstituted or substituted aryl, or

$R_7$  and  $R_8$ , together with the carbon atoms to which they are attached, form an unsaturated heterocycle having 5 or 6



ring atoms and one or two heteroatoms, independently of one another selected from nitrogen, oxygen and sulfur, which heterocycle may be substituted by one or two radicals which, independently of one another, are selected from the group consisting of C<sub>1</sub>-C<sub>4</sub>-alkyl, halogen, nitro, CN, halo-C<sub>1</sub>-C<sub>4</sub>-alkyl, OH, C<sub>1</sub>-C<sub>4</sub>-alkoxy, unsubstituted or substituted aryl, C<sub>2</sub>-C<sub>4</sub>-alkenyl, halo-C<sub>2</sub>-C<sub>4</sub>-alkenyl, C<sub>2</sub>-C<sub>4</sub>-alkynyl, halo-C<sub>2</sub>-C<sub>4</sub>-alkynyl;

R<sub>9</sub> is H, halogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>2</sub>-C<sub>6</sub>-haloalkenyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, C<sub>3</sub>-C<sub>6</sub>-halocycloalkyl, unsubstituted or substituted aryl;

R<sub>10</sub> is H, halogen, C<sub>1</sub>-C<sub>4</sub>-alkyl.

2. A compound of the formula 1 as claimed in claim 1 where:

W is -OCH<sub>2</sub>-, -C(R<sub>10</sub>)=N-O-CH<sub>2</sub>;

X is halogen, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy;

R<sub>1</sub> is H, C<sub>1</sub>-C<sub>4</sub>-alkyl, halogen, halo-C<sub>1</sub>-C<sub>4</sub>-alkyl;

R<sub>2</sub> is H, C<sub>1</sub>-C<sub>4</sub>-alkyl, halogen, halo-C<sub>1</sub>-C<sub>4</sub>-alkyl;

R<sub>3</sub> is H, C<sub>1</sub>-C<sub>4</sub>-alkyl;

n is 1 or 2;

R<sub>5</sub> is H or C<sub>1</sub>-C<sub>4</sub>-alkyl;

R<sub>6</sub> is H, C<sub>1</sub>-C<sub>4</sub>-alkyl, halo-C<sub>1</sub>-C<sub>4</sub>-alkyl;

R<sub>7</sub> is H, halogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, halo-C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, C<sub>3</sub>-C<sub>6</sub>-halocycloalkyl, phenyl;

R<sub>8</sub> is H, halogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, halo-C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, C<sub>3</sub>-C<sub>6</sub>-halocycloalkyl, C<sub>2</sub>-C<sub>6</sub>-alkenyl, phenyl which may be substituted by one or two halogen or C<sub>1</sub>-C<sub>4</sub>-alkyl; or

R<sub>7</sub> and R<sub>8</sub>, together with the carbon atoms to which they are attached, form an unsaturated heterocycle having 5 or 6 ring atoms and one or two heteroatoms, independently of one another selected from nitrogen, oxygen and sulfur, which heterocycle may be substituted by one or two

radicals which, independently of one another, are C<sub>1</sub>-C<sub>4</sub>-alkyl, halogen, halo-C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy and phenyl which may be substituted by one or two halogen or C<sub>1</sub>-C<sub>4</sub>-alkyl;

R<sub>9</sub> is H, halogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, halo-C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, C<sub>3</sub>-C<sub>6</sub>-halocycloalkyl, phenyl;

R<sub>10</sub> is H, halogen, C<sub>1</sub>-C<sub>4</sub>-alkyl.

3. A compound of the formula 1 as claimed in claim 1 where:

W is -OCH<sub>2</sub>-, -C(R<sub>10</sub>)=N-O-CH<sub>2</sub>;

X is halogen, C<sub>1</sub>-C<sub>4</sub>-alkoxy;

R<sub>1</sub> is H, C<sub>1</sub>-C<sub>4</sub>-alkyl;

R<sub>2</sub> is H, C<sub>1</sub>-C<sub>4</sub>-alkyl;

n is 1 or 2;

R<sub>3</sub> is C<sub>1</sub>-C<sub>4</sub>-alkyl;

R<sub>5</sub> is H, C<sub>1</sub>-C<sub>4</sub>-alkyl;

R<sub>6</sub> is H, C<sub>1</sub>-C<sub>4</sub>-alkyl;

R<sub>7</sub> is H, halogen, C<sub>1</sub>-C<sub>6</sub>-alkyl;

R<sub>8</sub> is H, halogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>2</sub>-C<sub>6</sub>-alkenyl; or

R<sub>7</sub> and R<sub>8</sub>, together with the carbon atoms to which they are attached, form a thiophenyl, furanyl, oxazolyl or thiazolyl radical, where these groups may have one or two substituents which are selected independently of one another from the group consisting of C<sub>1</sub>-C<sub>4</sub>-alkyl, halogen and phenyl which may be substituted by one or two halogen;

R<sub>9</sub> is H, halogen, C<sub>1</sub>-C<sub>6</sub>-alkyl;

R<sub>10</sub> is H, C<sub>1</sub>-C<sub>4</sub>-alkyl.

4. A compound of the formula 1 as claimed in claim 1 where:

W is  $-OCH_2-$ ,  $-C(R_{10})=N-O-CH_2$ ;

X is  $C_1-C_4$ -alkoxy;

$R_1$  is H;

$R_2$  is H,  $C_1-C_4$ -alkyl;

n is 1 or 2;

$R_3$  is  $C_1-C_4$ -alkyl;

$R_5$  is H,  $C_1-C_4$ -alkyl;

$R_6$  is H,  $C_1-C_4$ -alkyl;

$R_7$  is H, halogen;

$R_8$  is H,  $C_1-C_4$ -alkyl, halogen; or

$R_7$  and  $R_8$ , together with the carbon atoms to which they are attached, form a thiophenyl or oxazolyl radical, where these groups may be substituted by one or two halogen or phenyl and the phenyl may be substituted by one or two halogen;

$R_9$  is H, halogen;

$R_{10}$  is H,  $C_1-C_4$ -alkyl.

5. A compound of the formula 1 where:

W is  $-OCH_2-$ ,  $-C(R_{10})=N-O-CH_2-$ ;

X is halogen,  $C_1-C_4$ -alkyl,  $C_1-C_4$ -alkoxy;

$R_1$  is H,  $C_1-C_4$ -alkyl, halogen, nitro, CN, halo- $C_1-C_4$ -alkyl,  $C_1-C_4$ -alkoxy;

$R_2$  is H,  $C_1-C_4$ -alkyl, halogen, nitro, CN, halo- $C_1-C_4$ -alkyl,  $C_1-C_4$ -alkoxy;

n is 1 or 2;

$R_3$  is H,  $C_1-C_4$ -alkyl;

R<sub>5</sub> is H, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>2</sub>-C<sub>4</sub>-alkenyl;

R<sub>6</sub> is H, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkyl, C<sub>2</sub>-C<sub>4</sub>-alkenyl, aryl;

R<sub>7</sub> is H, halogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>2</sub>-C<sub>6</sub>-haloalkenyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, C<sub>3</sub>-C<sub>6</sub>-halocycloalkyl, unsubstituted or substituted aryl;


R<sub>8</sub> is H, halogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>2</sub>-C<sub>6</sub>-haloalkenyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, C<sub>3</sub>-C<sub>6</sub>-halocycloalkyl, unsubstituted or substituted aryl;

R<sub>9</sub> is H, halogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>2</sub>-C<sub>6</sub>-haloalkenyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, C<sub>3</sub>-C<sub>6</sub>-halocycloalkyl, unsubstituted or substituted aryl;

R<sub>10</sub> is H, halogen, C<sub>1</sub>-C<sub>4</sub>-alkyl.

6. The use of the compounds of the formula 1 as claimed in claim 1 as fungicides or for controlling pests.
7. A fungicide, comprising solid and/or liquid carriers and a fungicidally effective amount of at least one compound of the formula 1 as claimed in claim 1.
8. A method for controlling fungi, wherein the fungi or the materials, plants, seeds or the soil threatened by fungal attack is treated with a fungicidally effective amount of at least one compound of the formula 1 as claimed in claim 1.
9. A composition for controlling pests, comprising inert additives and a pesticidally effective amount of at least one compound of the formula 1 as claimed in claim 1.
10. A method for controlling pests, wherein the pests and/or their habitat are treated with a pesticidally effective amount of at least one compound of the formula 1 as claimed in claim 1.

11. A compound of the formula 6



in which

$W$ ,  $R_1$ ,  $R_2$ ,  $R_5$ ,  $R_6$ ,  $R_7$ ,  $R_8$ ,  $R_9$  and  $n$  are as defined in any of claims 1 to 5 and  $Y$  is  $NH_2$ .

10/010760


1

JC10 Rec'd PCT/PTO 12 MAR 2002

Unsaturated oxime ethers and the use thereof for control of harmful fungi and veterinary pests

- 5 The present invention relates to unsaturated oxime ethers, to intermediates for their preparation and to their use for controlling harmful fungi and animal pests.

Fungicidal oxime ether compounds are already known. WO 95/14009,  
 10 WO 96/17851, WO 96/36229, WO 96/36615, WO 96/36616, WO 96/36633, WO 97/0612, WO 98/05652 and WO 98/23155, for example, describe strobilurins having triazolone pharmacophores, of the formula:



in which  $R_{17}$  and  $R_{18}$  are H,  $C_1$ - $C_3$ -alkyl or phenyl.

- 25 Compounds having unsaturated oxime ether side-chains have already been disclosed in EP 386561A, EP 579124A, EP 585751A, EP 672347A, EP 673923A, WO 97/30032 and WO 97/33874.


It is an object of the present invention to provide compounds  
 30 having improved activity and/or a broader activity spectrum.

We have found that this object is achieved by compounds having a triazolone pharmacophore and an unsaturated oxime ether side-chain.

- 35 Accordingly, the present invention relates to unsaturated oxime ether compounds of the formula 1

40

45



10 where

W is  $-OCH_2-$ ,  $-C(R_{10})=N-O-CH_2-$  (where the  $CH_2$  end is in each case  
15 attached to the phenyl group carrying the triazolone radical);

X is halogen,  $C_1-C_4$ -alkyl,  $C_1-C_4$ -alkoxy;

$R_1$  is H,  $C_1-C_4$ -alkyl, halogen, nitro, CN, halo- $C_1-C_4$ -alkyl,  
20  $C_1-C_4$ -alkoxy;

$R_2$  is H,  $C_1-C_4$ -alkyl, halogen, nitro, CN, halo- $C_1-C_4$ -alkyl,  
 $C_1-C_4$ -alkoxy;

25 n is 1 or 2;

$R_3$  is H,  $C_1-C_4$ -alkyl;

$R_5$  is H,  $C_1-C_4$ -alkyl,  $C_2-C_4$ -alkenyl;

30  $R_6$  is H,  $C_1-C_4$ -alkyl,  $C_1-C_4$ -haloalkyl,  $C_2-C_4$ -alkenyl, aryl;

$R_7$  is H, halogen,  $C_1-C_6$ -alkyl,  $C_1-C_6$ -haloalkyl,  $C_2-C_6$ -alkenyl,  
35  $C_2-C_6$ -haloalkenyl,  $C_3-C_6$ -cycloalkyl,  $C_3-C_6$ -halocycloalkyl,  
unsubstituted or substituted aryl;

$R_8$  is H, halogen,  $C_1-C_6$ -alkyl,  $C_1-C_6$ -haloalkyl,  $C_2-C_6$ -alkenyl,  
 $C_2-C_6$ -haloalkenyl,  $C_3-C_6$ -cycloalkyl,  $C_3-C_6$ -halocycloalkyl,  
unsubstituted or substituted aryl, or

40  $R_7$  and  $R_8$ , together with the carbon atoms to which they are  
attached, form an unsaturated heterocycle having 5 or 6 ring  
atoms and one or two heteroatoms, independently of one  
another selected from nitrogen, oxygen and sulfur, which  
45 heterocycle may be substituted by one or two radicals which,  
independently of one another, are selected from the group  
consisting of  $C_1-C_4$ -alkyl, halogen, nitro, CN,

## 3

halo-C<sub>1</sub>-C<sub>4</sub>-alkyl, OH, C<sub>1</sub>-C<sub>4</sub>-alkoxy, unsubstituted or substituted aryl, C<sub>2</sub>-C<sub>4</sub>-alkenyl, halo-C<sub>2</sub>-C<sub>4</sub>-alkenyl, C<sub>2</sub>-C<sub>4</sub>-alkynyl, halo-C<sub>2</sub>-C<sub>4</sub>-alkynyl;

- 5 R<sub>9</sub> is H, halogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>2</sub>-C<sub>6</sub>-haloalkenyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, C<sub>3</sub>-C<sub>6</sub>-halocycloalkyl, unsubstituted or substituted aryl;

R<sub>10</sub> is H, halogen, C<sub>1</sub>-C<sub>4</sub>-alkyl.

10

The meanings listed above are collective terms for individual enumerations of the individual group members. All hydrocarbon chains can be straight-chain or branched. Halogenated substituents preferably carry 1 to 5 identical or different

15 halogen atoms.

Specific meanings are, for example:

- halogen: fluorine, chlorine, bromine, iodine, preferably  
20 fluorine or chlorine;
- C<sub>1</sub>-C<sub>4</sub>-alkyl: methyl, ethyl, n-propyl, 1-methylethyl, n-butyl, 1-methylpropyl, 2-methylpropyl, 1,1-dimethylethyl, preferably methyl;
- 25 - C<sub>1</sub>-C<sub>6</sub>-alkyl: C<sub>1</sub>-C<sub>4</sub>-alkyl as mentioned above, and also n-pentyl, 1-methylbutyl, 2-methylbutyl, 3-methylbutyl, 2,2-dimethylpropyl, 1-ethylpropyl, n-hexyl, 1,1-dimethylpropyl, 1,2-dimethylpropyl, 1-methylpentyl,  
30 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 2,2-dimethylbutyl, 2,3-dimethylbutyl, 3,3-dimethylbutyl, 1-ethylbutyl, 2-ethylbutyl, 1,1,2-trimethylpropyl, 1,2,2-trimethylpropyl, 1-ethyl-1-methylpropyl,  
35 1-ethyl-2-methylpropyl, preferably methyl, ethyl, n-propyl, 1-methylethyl, n-butyl or 1,1-dimethylethyl;
- C<sub>2</sub>-C<sub>6</sub>-alkenyl: ethenyl, prop-1-en-1-yl, prop-2-en-1-yl, 1-methylethenyl, n-buten-1-yl, n-buten-2-yl, n-buten-3-yl,  
40 1-methylprop-1-en-1-yl, 2-methylprop-1-en-1-yl, 1-methylprop-2-en-1-yl, 2-methylprop-2-en-1-yl, n-penten-1-yl, n-penten-2-yl, n-penten-3-yl, n-penten-4-yl, 1-methylbut-1-en-1-yl, 2-methylbut-1-en-1-yl, 3-methylbut-1-en-1-yl, 1-methylbut-2-en-1-yl,  
45 2-methylbut-2-en-1-yl, 3-methylbut-2-en-1-yl, 1-methylbut-3-en-1-yl, 2-methylbut-3-en-1-yl, 3-methylbut-3-en-1-yl, 1,1-dimethylprop-2-en-1-yl,



- 1,2-dimethylprop-1-en-1-yl, 1,2-dimethylprop-2-en-1-yl,  
1-ethylprop-1-en-2-yl, 1-ethylprop-2-en-1-yl,  
n-hex-1-en-1-yl, n-hex-2-en-1-yl, n-hex-3-en-1-yl,  
n-hex-4-en-1-yl, n-hex-5-en-1-yl, 1-methylpent-1-en-1-yl,  
5 2-methylpent-1-en-1-yl, 3-methylpent-1-en-1-yl,  
4-methylpent-1-en-1-yl, 1-methylpent-2-en-1-yl,  
2-methylpent-2-en-1-yl, 3-methylpent-2-en-1-yl,  
4-methylpent-2-en-1-yl, 1-methylpent-3-en-1-yl,  
2-methylpent-3-en-1-yl, 3-methylpent-3-en-1-yl,  
10 4-methylpent-3-en-1-yl, 1-methylpent-4-en-1-yl,  
2-methylpent-4-en-1-yl, 3-methylpent-4-en-1-yl,  
4-methylpent-4-en-1-yl, 1,1-dimethylbut-2-en-1-yl,  
1,1-dimethylbut-3-en-1-yl, 1,2-dimethylbut-1-en-1-yl,  
1,2-dimethylbut-2-en-1-yl, 1,2-dimethylbut-3-en-1-yl,  
15 1,3-dimethylbut-1-en-1-yl, 1,3-dimethylbut-2-en-1-yl,  
1,3-dimethylbut-3-en-1-yl, 2,2-dimethylbut-3-en-1-yl,  
2,3-dimethylbut-1-en-1-yl, 2,3-dimethylbut-2-en-1-yl,  
2,3-dimethylbut-3-en-1-yl, 3,3-dimethylbut-1-en-1-yl,  
3,3-dimethylbut-2-en-1-yl, 1-ethylbut-1-en-1-yl,  
20 1-ethylbut-2-en-1-yl, 1-ethylbut-3-en-1-yl,  
2-ethylbut-1-en-1-yl, 2-ethylbut-2-en-1-yl,  
2-ethylbut-3-en-1-yl, 1,1,2-trimethylprop-2-en-1-yl,  
1-ethyl-1-methylprop-2-en-1-yl,  
1-ethyl-2-methylprop-1-en-1-yl,  
25 1-ethyl-2-methylprop-2-en-1-yl, preferably ethenyl or  
prop-2-en-1-yl;
- C<sub>2</sub>-C<sub>4</sub>-alkynyl: ethynyl, prop-1-yn-1-yl, prop-2-yn-3-yl,  
n-but-1-yn-1-yl, n-but-1-yn-4-yl, n-but-2-yn-1-yl, preferably  
30 prop-2-yn-1-yl;
- C<sub>1</sub>-C<sub>2</sub>-haloalkyl: for example chloromethyl, dichloromethyl,  
trichloromethyl, fluoromethyl, difluoromethyl,  
trifluoromethyl, chlorofluoromethyl, dichlorofluoromethyl,  
35 chlorodifluoromethyl, 1-fluoroethyl, 2-fluoroethyl,  
2,2-difluoroethyl, 2,2,2-trifluoroethyl,  
2-chloro-2-fluoroethyl, 2-chloro-2,2-difluoroethyl,  
2,2-dichloro-2-fluoroethyl, 2,2,2-trichloroethyl,  
pentafluoroethyl, preferably difluoromethyl or  
40 trifluoromethyl;
- C<sub>1</sub>-C<sub>6</sub>-haloalkyl: C<sub>1</sub>-C<sub>6</sub>-alkyl as mentioned above which is  
partially or fully substituted by fluorine, chlorine and/or  
bromine, i.e., for example, the abovementioned C<sub>1</sub>-C<sub>2</sub>-haloalkyl  
45 radicals, and also 3-chloropropyl or heptafluoropropyl,

## 5

preferably trifluoromethyl, pentafluoroethyl or heptafluoropropyl;

- C<sub>2</sub>-C<sub>6</sub>-haloalkenyl: C<sub>2</sub>-C<sub>6</sub>-alkenyl as mentioned above which is partially or fully substituted by fluorine, chlorine and/or bromine, i.e., for example, 2-chloroallyl, 3-chloroallyl or 3,3-dichloroallyl;
  - C<sub>2</sub>-C<sub>4</sub>-haloalkynyl: C<sub>2</sub>-C<sub>4</sub>-alkynyl as mentioned above which is partially or fully substituted by fluorine, chlorine and/or bromine, for example chloroethynyl, 3-chloropropynyl;
  - C<sub>3</sub>-C<sub>6</sub>-cycloalkyl: cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, preferably cyclopropyl, cyclopentyl or cyclohexyl;
  - halo-C<sub>3</sub>-C<sub>6</sub>-cycloalkyl: C<sub>3</sub>-C<sub>6</sub>-cycloalkyl as mentioned above which is partially or fully substituted by fluorine, chlorine and/or bromine, i.e., for example, 2-, 3- or 4-chlorocyclopentyl, 2-, 3- or 4-chlorocyclohexyl, 2,3,4-trichlorocyclopentyl or 2,3,4,5,6-pentachlorocyclohexyl;
  - aryl is preferably phenyl or naphthyl, in particular phenyl.
- 25 The unsaturated heterocycle having 5 or 6 ring atoms can be aromatic or nonaromatic. Aromatic heterocycles are, in particular, 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, 2-pyrrolyl, 3-pyrrolyl, 3-isoxazolyl, 4-isoxazolyl, 5-isoxazolyl, 3-isothiazolyl, 4-isothiazolyl, 5-isothiazolyl, 3-pyrazolyl, 4-pyrazolyl, 5-pyrazolyl, 2-oxazolyl, 4-oxazolyl, 5-oxazolyl, 2-thiazolyl, 4-thiazolyl, 5-thiazolyl, 2-imidazolyl, 4-imidazolyl, 1,2,4-oxadiazol-3-yl, 1,2,4-oxadiazol-5-yl, 1,2,4-thiadiazol-3-yl, 1,2,4-thiadiazol-5-yl, 1,2,4-triazol-3-yl, 1,3,4-oxadiazol-2-yl, 1,3,4-thiadiazol-2-yl, 1,3,4-triazol-2-yl, 2-pyridinyl, 3-pyridinyl, 4-pyridinyl, 3-pyridazinyl, 4-pyridazinyl, 2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl, 2-pyrazinyl, in particular furanyl, thienyl, oxazolyl and thiazolyl.
- 40 If the aryl radical is substituted, it has preferably one or two substituents which are independently of one another selected from the group consisting of halogen, in particular fluorine or chlorine, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, nitro, OH and CN. Preference is given to halogen and/or C<sub>1</sub>-C<sub>4</sub>-alkyl.

## 6

The substituents in the formula 1 preferably have the following meaning:

- 5 X is C<sub>1</sub>-C<sub>4</sub>-alkoxy or halogen, in particular C<sub>1</sub>-C<sub>4</sub>-alkoxy;
- R<sub>1</sub> is H, C<sub>1</sub>-C<sub>4</sub>-alkyl, halogen, halo-C<sub>1</sub>-C<sub>4</sub>-alkyl, in particular H or C<sub>1</sub>-C<sub>4</sub>-alkyl;
- 10 R<sub>2</sub> is H, C<sub>1</sub>-C<sub>4</sub>-alkyl, halogen, halo-C<sub>1</sub>-C<sub>4</sub>-alkyl, in particular H or C<sub>1</sub>-C<sub>4</sub>-alkyl;
- R<sub>3</sub> is C<sub>1</sub>-C<sub>4</sub>-alkyl;
- 15 R<sub>5</sub> is H or C<sub>1</sub>-C<sub>4</sub>-alkyl;
- R<sub>6</sub> is H, C<sub>1</sub>-C<sub>4</sub>-alkyl or halo-C<sub>1</sub>-C<sub>4</sub>-alkyl, in particular H or C<sub>1</sub>-C<sub>4</sub>-alkyl;
- 20 R<sub>7</sub> is H, halogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, halo-C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, C<sub>3</sub>-C<sub>6</sub>-halocycloalkyl or phenyl, in particular H, halogen, C<sub>1</sub>-C<sub>6</sub>-alkyl and particularly preferably H or halogen;
- 25 R<sub>8</sub> is H, halogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, halo-C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, C<sub>3</sub>-C<sub>6</sub>-halocycloalkyl, C<sub>2</sub>-C<sub>6</sub>-alkenyl or phenyl, in particular H, halogen, C<sub>1</sub>-C<sub>6</sub>-alkyl; or
- 30 R<sub>7</sub> and R<sub>8</sub>, together with the carbon atoms to which they are attached, are a thienyl, furyl, oxazolyl or thiazolyl radical, where these groups may be substituted by C<sub>1</sub>-C<sub>4</sub>-alkyl, halogen or phenyl which may be substituted by one or two halogen, and are in particular thienyl or oxazolyl, where these groups may be substituted by one or two halogen or phenyl and the phenyl substituent for its
- 35 part may be substituted by one or two halogen;
- R<sub>9</sub> is H, halogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, halo-C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, C<sub>3</sub>-C<sub>6</sub>-halocycloalkyl or phenyl, in particular H, halogen, C<sub>1</sub>-C<sub>6</sub>-alkyl and particularly
- 40 preferably H or halogen;
- R<sub>10</sub> is H or C<sub>1</sub>-C<sub>4</sub>-alkyl.

The oxime side-chain may be attached to the phenyl ring in the

45 o-, m- or p-position to W. Preference is given to the p-position.

The radical  $R_1$  is preferably located in the 6-position.

If the oxime side-chain is attached in the p-position to W, the radical  $R_2$  is preferably in the 2- and/or 5-position.

5

Preferred embodiments are the compounds of the formula 1 where:

A)

10 W is  $-OCH_2-$ ,  $-C(R_{10})=N-O-CH_2$ ;

X is halogen,  $C_1-C_4$ -alkyl,  $C_1-C_4$ -alkoxy;

$R_1$  is H,  $C_1-C_4$ -alkyl, halogen, halo- $C_1-C_4$ -alkyl;

15

$R_2$  is H,  $C_1-C_4$ -alkyl, halogen, halo- $C_1-C_4$ -alkyl;

$R_3$  is H,  $C_1-C_4$ -alkyl;

20 n is 1 or 2;

$R_5$  is H or  $C_1-C_4$ -alkyl;

$R_6$  is H,  $C_1-C_4$ -alkyl, halo- $C_1-C_4$ -alkyl;

25

$R_7$  is H, halogen,  $C_1-C_6$ -alkyl, halo- $C_1-C_6$ -alkyl,  $C_3-C_6$ -cycloalkyl,  $C_3-C_6$ -halocycloalkyl, phenyl;

30

$R_8$  is H, halogen,  $C_1-C_6$ -alkyl, halo- $C_1-C_6$ -alkyl,  $C_3-C_6$ -cycloalkyl,  $C_3-C_6$ -halocycloalkyl,  $C_2-C_6$ -alkenyl, phenyl; or

35

$R_7$  and  $R_8$ , together with the carbon atoms to which they are attached, form an unsaturated heterocycle having 5 or 6 ring atoms and one or two heteroatoms, independently of one another selected from nitrogen, oxygen and sulfur, which heterocycle may be substituted by one or two radicals which, independently of one another, are  $C_1-C_4$ -alkyl, halogen, halo- $C_1-C_4$ -alkyl,  $C_1-C_4$ -alkoxy and phenyl which may be substituted by one or two halogen or  $C_1-C_4$ -alkyl;

40

$R_9$  is H, halogen,  $C_1-C_6$ -alkyl, halo- $C_1-C_6$ -alkyl,  $C_3-C_6$ -cycloalkyl,  $C_3-C_6$ -halocycloalkyl, phenyl;

45

$R_{10}$  is H, halogen,  $C_1-C_4$ -alkyl;

B)

W is  $-\text{OCH}_2-$ ,  $-\text{C}(\text{R}_{10})=\text{N}-\text{O}-\text{CH}_2$ ;5 X is halogen,  $\text{C}_1$ - $\text{C}_4$ -alkoxy; $\text{R}_1$  is H,  $\text{C}_1$ - $\text{C}_4$ -alkyl; $\text{R}_2$  is H,  $\text{C}_1$ - $\text{C}_4$ -alkyl;

10

n is 1 or 2;

 $\text{R}_3$  is  $\text{C}_1$ - $\text{C}_4$ -alkyl;15  $\text{R}_5$  is H,  $\text{C}_1$ - $\text{C}_4$ -alkyl; $\text{R}_6$  is H,  $\text{C}_1$ - $\text{C}_4$ -alkyl; $\text{R}_7$  is H, halogen,  $\text{C}_1$ - $\text{C}_6$ -alkyl;

20

 $\text{R}_8$  is H, halogen,  $\text{C}_1$ - $\text{C}_6$ -alkyl; or

$\text{R}_7$  and  $\text{R}_8$ , together with the carbon atoms to which they are attached, form a thienyl, furanyl, oxazolyl or thiazolyl radical, where these groups may have one or two substituents which are selected independently of one another from the group consisting of  $\text{C}_1$ - $\text{C}_4$ -alkyl, halogen and phenyl which may be substituted by one or two halogen;

30

 $\text{R}_9$  is H, halogen,  $\text{C}_1$ - $\text{C}_6$ -alkyl; $\text{R}_{10}$  is H,  $\text{C}_1$ - $\text{C}_4$ -alkyl;

35 C)

W is  $-\text{OCH}_2-$ ,  $-\text{C}(\text{R}_{10})=\text{N}-\text{O}-\text{CH}_2$ ;X is  $\text{C}_1$ - $\text{C}_4$ -alkoxy;

40

 $\text{R}_1$  is H; $\text{R}_2$  is H,  $\text{C}_1$ - $\text{C}_4$ -alkyl;

45

n is 1 or 2;

R<sub>5</sub> is H, C<sub>1</sub>-C<sub>4</sub>-alkyl;

R<sub>7</sub> is H, halogen;

10


15

R<sub>10</sub> is H, C<sub>1</sub>-C<sub>4</sub>-alkyl;

20

D) compounds of the formula Ia:

25



35 where

R<sub>2</sub> is chlorine, methyl;

40

R<sub>5</sub> is H, methyl, ethyl;

45

R<sub>7</sub> is H, methyl, chlorine;

10

$R_8$  is H, methyl, ethyl, chlorine, phenyl, ethenyl, prop-1-en-1-yl;


$R_9$  is H, methyl, chlorine;

5

E) compounds of the formula 1b:

10

15



1b

20 where:

X is chlorine, methoxy;

25  $R_5$  is H, methyl, ethyl;

$R_7$  is H, methyl, chlorine;

30  $R_8$  is H, methyl, ethyl, chlorine, phenyl, ethenyl, prop-1-en-1-yl;


$R_9$  is H, methyl, chlorine.

The oxime side-chain is preferably attached in the p-position.

35 F) Compounds of the formula 1c:

40

45



1c

in which R is:

thiophen-2-yl, thiophen-3-yl, furan-2-yl, furan-3-yl,  
5-chlorothiophen-2-yl, 5-bromothiophen-2-yl, 5-chlorofuran-2-yl,  
5 5-bromofuran-2-yl, 5-methylthiophen-2-yl, 5-phenylthiophen-2-yl,  
5-methylfuran-2-yl, 5-phenylfuran-2-yl, oxazol-4-yl, oxazol-5-yl,  
2-(p-chlorophenyl)oxazol-4-yl, 2-(p-bromophenyl)oxazol-4-yl,  
thiazol-4-yl, thiazol-5-yl.

10 Here, the radicals X, R<sub>2</sub> and R<sub>5</sub> have the meanings mentioned above or, preferably, the following meanings:

X is chlorine, methoxy;

15 R<sub>2</sub> is 2-methyl, 2-chloro, 2,5-dimethyl;


R<sub>5</sub> is H, methyl, ethyl.

G) Compounds of the formulae 1d, 1e and 1f:

20


25

30




35

40



45





5 in which R is:

thiophen-2-yl, thiophen-3-yl, furan-2-yl, furan-3-yl,  
 5-chlorothiophen-2-yl, 5-bromothiophen-2-yl, 5-chlorofuran-2-yl,  
 5-bromofuran-2-yl, 5-methylthiophen-2-yl, 5-phenylthiophen-2-yl,  
 15 5-methylfuran-2-yl, 5-phenylfuran-2-yl, oxazol-4-yl, oxazol-5-yl,  
 2-(p-chlorophenyl)oxazol-4-yl, 2-(p-bromophenyl)oxazol-4-yl,  
 thiazol-4-yl, thiazol-5-yl.

Here, the radicals X and R<sub>5</sub> have the meanings given above or,  
 20 preferably, the following meanings:

X is chlorine, methoxy;

R<sub>5</sub> is H, methyl, ethyl.

25 In particular with respect to their use, preference is given to  
 the compounds of the formulae 1c to 1f compiled below. The groups  
 mentioned here for a substituent are furthermore on their own,  
 independently of the combination in which they are mentioned, a  
 30 particularly preferred embodiment of the respective substituent:

- a) Compounds of the formula 1c in which R<sub>2</sub> is 2-methyl (based on  
 the oxygen substituent), X is chlorine and R<sub>5</sub> is hydrogen and  
 R corresponds to a row of Table 1.
- 35 b) Compounds of the formula 1c in which R<sub>2</sub> is 2-methyl (based on  
 the oxygen substituent), X is methoxy and R<sub>5</sub> is hydrogen and  
 R corresponds to a row of Table 1.
- 40 c) Compounds of the formula 1c in which R<sub>2</sub> is 2-chloro (based on  
 the oxygen substituent), X is chlorine and R<sub>5</sub> is hydrogen and  
 R corresponds to a row of Table 1.
- d) Compounds of the formula 1c in which R<sub>2</sub> is 2-chloro (based on  
 45 the oxygen substituent), X is methoxy and R<sub>5</sub> is hydrogen and  
 R corresponds to a row of Table 1.

## 13

- e) Compounds of the formula 1c in which  $R_2$  is 2,5-dimethyl (based on the oxygen substituent), X is chlorine and  $R_5$  is hydrogen and R corresponds to a row of Table 1.
- 5 f) Compounds of the formula 1c in which  $R_2$  is 2,5-dimethyl (based on the oxygen substituent), X is methoxy and  $R_5$  is hydrogen and R corresponds to a row of Table 1.
- 10 g) Compounds of the formula 1c in which  $R_2$  is 2-methyl (based on the oxygen substituent), X is chlorine and  $R_5$  is methyl and R corresponds to a row of Table 1.
- 15 h) Compounds of the formula 1c in which  $R_2$  is 2-methyl (based on the oxygen substituent), X is methoxy and  $R_5$  is methyl and R corresponds to a row of Table 1.
- 20 i) Compounds of the formula 1c in which  $R_2$  is 2-chloro (based on the oxygen substituent), X is chlorine and  $R_5$  is methyl and R corresponds to a row of Table 1.
- j) Compounds of the formula 1c in which  $R_2$  is 2-chloro (based on the oxygen substituent), X is methoxy and  $R_5$  is methyl and R corresponds to a row of Table 1.
- 25 k) Compounds of the formula 1c in which  $R_2$  is 2,5-dimethyl (based on the oxygen substituent), X is chlorine and  $R_5$  is methyl and R corresponds to a row of Table 1.
- 30 l) Compounds of the formula 1c in which  $R_2$  is 2,5-dimethyl (based on the oxygen substituent), X is methoxy and  $R_5$  is methyl and R corresponds to a row of Table 1.
- 35 m) Compounds of the formula 1c in which  $R_2$  is 2-methyl (based on the oxygen substituent), X is chlorine and  $R_5$  is ethyl and R corresponds to a row of Table 1.
- 40 n) Compounds of the formula 1c in which  $R_2$  is 2-methyl (based on the oxygen substituent), X is methoxy and  $R_5$  is ethyl and R corresponds to a row of Table 1.
- o) Compounds of the formula 1c in which  $R_2$  is 2-chloro (based on the oxygen substituent), X is chlorine and  $R_5$  is ethyl and R corresponds to a row of Table 1.

## 14

- p) Compounds of the formula 1c in which  $R_2$  is 2-chloro (based on the oxygen substituent), X is methoxy and  $R_5$  is ethyl and R corresponds to a row of Table 1.
- 5 q) Compounds of the formula 1c in which  $R_2$  is 2,5-dimethyl (based on the oxygen substituent), X is chlorine and  $R_5$  is ethyl and R corresponds to a row of Table 1.
- 10 r) Compounds of the formula 1c in which  $R_2$  is 2,5-dimethyl (based on the oxygen substituent), X is methoxy and  $R_5$  is ethyl and R corresponds to a row of Table 1.
- s) Compounds of the formula 1d in which X is chlorine and  $R_5$  is hydrogen and R corresponds to a row of Table 1.
- 15 t) Compounds of the formula 1d in which X is methoxy and  $R_5$  is hydrogen and R corresponds to a row of Table 1.
- 20 u) Compounds of the formula 1d in which X is chlorine and  $R_5$  is methyl and R corresponds to a row of Table 1.
- v) Compounds of the formula 1d in which X is methoxy and  $R_5$  is methyl and R corresponds to a row of Table 1.
- 25 w) Compounds of the formula 1d in which X is chlorine and  $R_5$  is ethyl and R corresponds to a row of Table 1.
- x) Compounds of the formula 1d in which X is methoxy and  $R_5$  is ethyl and R corresponds to a row of Table 1.
- 30 y) Compounds of the formula 1e in which X is chlorine and  $R_5$  is hydrogen and R corresponds to a row of Table 1.
- 35 z) Compounds of the formula 1e in which X is methoxy and  $R_5$  is hydrogen and R corresponds to a row of Table 1.
- aa) Compounds of the formula 1e in which X is chlorine and  $R_5$  is methyl and R corresponds to a row of Table 1.
- 40 ab) Compounds of the formula 1e in which X is methoxy and  $R_5$  is methyl and R corresponds to a row of Table 1.
- ac) Compounds of the formula 1e in which X is chlorine and  $R_5$  is ethyl and R corresponds to a row of Table 1.

## 15

- ad) Compounds of the formula 1e in which X is methoxy and R<sub>5</sub> is ethyl and R corresponds to a row of Table 1.
- ae) Compounds of the formula 1f in which X is chlorine and R<sub>5</sub> is hydrogen and R corresponds to a row of Table 1.
- af) Compounds of the formula 1f in which X is methoxy and R<sub>5</sub> is hydrogen and R corresponds to a row of Table 1.
- 10 ag) Compounds of the formula 1f in which X is chlorine and R<sub>5</sub> is methyl and R corresponds to a row of Table 1.
- ah) Compounds of the formula 1f in which X is methoxy and R<sub>5</sub> is methyl and R corresponds to a row of Table 1.
- 15 ai) Compounds of the formula 1f in which X is chlorine and R<sub>5</sub> is ethyl and R corresponds to a row of Table 1.
- aj) Compounds of the formula 1f in which X is methoxy and R<sub>5</sub> is ethyl and R corresponds to a row of Table 1.
- 20

Table 1

(E), (E,E) and (Z) refer to the substituents on the double bond indicated in R.

25

Number	R
1.	-CH=CH <sub>2</sub>
2.	(E) -CH=CH-CH <sub>3</sub>
30 3.	(Z) -CH=CH-CH <sub>3</sub>
4.	-CH=C(CH <sub>3</sub> ) <sub>2</sub>
5.	(E) -CH=CH-C <sub>2</sub> H <sub>5</sub>
6.	(Z) -CH=CH-C <sub>2</sub> H <sub>5</sub>
7.	(E) -CH=C(CH <sub>3</sub> )-C <sub>2</sub> H <sub>5</sub>
8.	(Z) -CH=C(CH <sub>3</sub> )-C <sub>2</sub> H <sub>5</sub>
35 9.	-CH=C(C <sub>2</sub> H <sub>5</sub> ) <sub>2</sub>
10.	(E) -CH=CH-Cl
11.	(Z) -CH=CH-Cl <sub>3</sub>
12.	(E) -CH=C(Cl)-CH
13.	(Z) -CH=C(Cl)-CH <sub>3</sub>
40 14.	(E) -CH=C(Cl)-C <sub>2</sub> H <sub>5</sub>
15.	(Z) -CH=C(Cl)-C <sub>2</sub> H <sub>5</sub>
16.	-CH=CCl <sub>2</sub>
17.	(E) -CH=CH-CH=CH <sub>2</sub>
18.	(E,E) -CH=CH-CH=CH-CH <sub>3</sub>
19.	(E,E) -CH=CH-CH=CH-C <sub>6</sub> H <sub>5</sub>
45 20.	(E,E) -CH=CH-CH=CH-(p-F-C <sub>6</sub> H <sub>4</sub> )
21.	(E,E) -CH=CH-CH=CH-(p-Cl-C <sub>6</sub> H <sub>4</sub> )
22.	-C(CH <sub>3</sub> )=CH <sub>2</sub>

	23.	(E)	$-\text{C}(\text{CH}_3)=\text{CH}-\text{CH}_3$
	24.	(Z)	$-\text{C}(\text{CH}_3)=\text{CH}-\text{CH}_3$
	25.		$-\text{C}(\text{CH}_3)=\text{C}(\text{CH}_3)_2$
	26.	(E)	$-\text{C}(\text{CH}_3)=\text{CH}-\text{C}_2\text{H}_5$
5	27.	(Z)	$-\text{C}(\text{CH}_3)=\text{CH}-\text{C}_2\text{H}_5$
	28.	(E)	$-\text{C}(\text{CH}_3)=\text{C}(\text{CH}_3)-\text{C}_2\text{H}_5$
	29.	(Z)	$-\text{C}(\text{CH}_3)=\text{C}(\text{CH}_3)-\text{C}_2\text{H}_5$
	30.		$-\text{C}(\text{CH}_3)=\text{C}(\text{C}_2\text{H}_5)_2$
	31.	(E)	$-\text{C}(\text{CH}_3)=\text{CH}-\text{Cl}$
	32.	(Z)	$-\text{C}(\text{CH}_3)=\text{CH}-\text{Cl}$
10	33.	(E)	$-\text{C}(\text{CH}_3)=\text{C}(\text{Cl})-\text{CH}_3$
	34.	(Z)	$-\text{C}(\text{CH}_3)=\text{C}(\text{Cl})-\text{CH}_3$
	35.	(E)	$-\text{C}(\text{CH}_3)=\text{C}(\text{Cl})-\text{C}_2\text{H}_5$
	36.	(Z)	$-\text{C}(\text{CH}_3)=\text{C}(\text{Cl})-\text{C}_2\text{H}_5$
	37.		$-\text{C}(\text{CH}_3)=\text{CCl}_2$
15	38.	(E)	$-\text{C}(\text{CH}_3)=\text{CH}-\text{CH}=\text{CH}_2$
	39.	(E,E)	$-\text{C}(\text{CH}_3)=\text{CH}-\text{CH}=\text{CH}-\text{CH}_3$
	40.	(E,E)	$-\text{C}(\text{CH}_3)=\text{CH}-\text{CH}=\text{CH}-\text{C}_6\text{H}_5$
	41.	(E,E)	$-\text{C}(\text{CH}_3)=\text{CH}-\text{CH}=\text{CH}-(p\text{-F}-\text{C}_6\text{H}_4)$
	42.	(E,E)	$-\text{C}(\text{CH}_3)=\text{CH}-\text{CH}=\text{CH}-(p\text{-Cl}-\text{C}_6\text{H}_4)$
	43.		$-\text{C}(\text{C}_2\text{H}_5)=\text{CH}_2$
20	44.	(E)	$-\text{C}(\text{C}_2\text{H}_5)=\text{CH}-\text{CH}_3$
	45.	(Z)	$-\text{C}(\text{C}_2\text{H}_5)=\text{CH}-\text{CH}_3$
	46.		$-\text{C}(\text{C}_2\text{H}_5)=\text{C}(\text{CH}_3)_2$
	47.	(E)	$-\text{C}(\text{C}_2\text{H}_5)=\text{CH}-\text{C}_2\text{H}_5$
	48.	(Z)	$-\text{C}(\text{C}_2\text{H}_5)=\text{CH}-\text{C}_2\text{H}_5$
25	49.	(E)	$-\text{C}(\text{C}_2\text{H}_5)=\text{C}(\text{CH}_3)-\text{C}_2\text{H}_5$
	50.	(Z)	$-\text{C}(\text{C}_2\text{H}_5)=\text{C}(\text{CH}_3)-\text{C}_2\text{H}_5$
	51.		$-\text{C}(\text{C}_2\text{H}_5)=\text{C}(\text{C}_2\text{H}_5)_2$
	52.	(E)	$-\text{C}(\text{C}_2\text{H}_5)=\text{CH}-\text{Cl}$
	53.	(Z)	$-\text{C}(\text{C}_2\text{H}_5)=\text{CH}-\text{Cl}$
	54.	(E)	$-\text{C}(\text{C}_2\text{H}_5)=\text{C}(\text{Cl})-\text{CH}_3$
30	55.	(Z)	$-\text{C}(\text{C}_2\text{H}_5)=\text{C}(\text{Cl})-\text{CH}_3$
	56.	(E)	$-\text{C}(\text{C}_2\text{H}_5)=\text{C}(\text{Cl})-\text{C}_2\text{H}_5$
	57.	(Z)	$-\text{C}(\text{C}_2\text{H}_5)=\text{C}(\text{Cl})-\text{C}_2\text{H}_5$
	58.		$-\text{C}(\text{C}_2\text{H}_5)=\text{CCl}_2$
	59.	(E)	$-\text{C}(\text{C}_2\text{H}_5)=\text{CH}-\text{CH}=\text{CH}_2$
35	60.	(E,E)	$-\text{C}(\text{C}_2\text{H}_5)=\text{CH}-\text{CH}=\text{CH}-\text{CH}_3$
	61.	(E,E)	$-\text{C}(\text{C}_2\text{H}_5)=\text{CH}-\text{CH}=\text{CH}-\text{C}_6\text{H}_5$
	62.	(E,E)	$-\text{C}(\text{C}_2\text{H}_5)=\text{CH}-\text{CH}=\text{CH}-(p\text{-F}-\text{C}_6\text{H}_4)$
	63.	(E,E)	$-\text{C}(\text{C}_2\text{H}_5)=\text{CH}-\text{CH}=\text{CH}-(p\text{-Cl}-\text{C}_6\text{H}_4)$
	64.		thiophen-2-yl
	65.		thiophen-3-yl
40	66.		furan-2-yl
	67.		furan-3-yl
	68.		5-chlorothiophen-2-yl
	69.		5-bromothiophen-2-yl
	70.		5-chlorofuran-2-yl
45	71.		5-bromofuran-2-yl
	72.		5-methylthiophen-2-yl
	73.		5-phenylthiophen-2-yl
	74.		5-methylfuran-2-yl


5	75.	5-phenylfuran-2-yl
	76.	oxazol-4-yl
	77.	oxazol-5-yl
	78.	thiazol-4-yl
	79.	thiazol-5-yl
10	80.	2-phenyloxazol-4-yl
	81.	2-phenylthiazol-4-yl
	82.	2-(p-chlorophenyl)oxazol-4-yl
	83.	2-(p-chlorophenyl)thiazol-4-yl
	84.	2-(p-bromophenyl)oxazol-4-yl
	85.	2-(p-bromophenyl)thiazol-4-yl
	86.	2-(p-fluorophenyl)oxazol-4-yl
	87.	2-(p-fluorophenyl)thiazol-4-yl
	88.	2-(2,4-dichlorophenyl)oxazol-4-yl
	89.	2-(2,4-dichlorophenyl)thiazol-4-yl

- 15 The compounds according to the invention can be prepared by a plurality of methods. The preparation is preferably carried out by one of the synthesis routes A or B which are illustrated in the formula scheme below. The synthesis starts with a compound of
- 20 the formula 2 in which  $X_1$  in the case of route A is  $CH_3$  or  $PGOCH_2-$ , where PG is a protective group for benzyl ethers. After the synthesis of the triazolone ring, the protective group is cleaved off and the resulting benzyl alcohol is converted by methods known from the literature into  $X_2CH_2-$ ,  $X_2$  being a
- 25 nucleophilically removable group (cf. T. W. Greene: "Protective Groups in Organic Synthesis", 2nd Edition 1991, John Wiley, New York). In the case of route B,  $X_1 = X_2CH_2-$ , where  $X_2$  is a nucleophilically removable group, such as, for example, chlorine, fluorine, bromine, nitro, alkyl- or arylsulfonate, such as
- 30 mesylate, tosylate or triflate. The preparation of these compounds is known and, for example for the synthesis from the corresponding nitro compounds, described in Houben-Weyl, Vol. IV/1c, 4th Edition, p. 506ff, Thieme Verlag, Stuttgart 1980.

35

40

45



The amino function of the compounds 2 is initially converted into a thiazolone group, giving the compounds 3. The reaction sequence which leads to this conversion is described in detail in

- 5 WO 95/14009 (compounds Nos. 21  $\rightarrow$  19  $\rightarrow$  17  $\rightarrow$  1), and also in WO 96/26191 and DE 198 09 995.

The group  $X_1$  is subsequently converted into a reactive group which can react with the radical V in the compound 7 or the radical  $W_1$

- 10 in the compound of the formula 8, linking the two phenyl rings and forming the group W. The group V is a phenolic OH group or the group  $C(R_{10})=N-OH$ . Thus, a leaving group is introduced into the group  $X_1$  and removed with the hydrogen atom of the phenolic or oximic OH group in the presence of a base. Examples of such
- 15 leaving groups are chloride, bromide, p-toluenesulfonate, methanesulfonate or trifluoromethanesulfonate. These leaving groups are introduced in a manner known to the person skilled in the art, for example by reacting the compound 3 where  $X_1 = CH_3$  with N-bromosuccinimide or N-chlorosuccinimide.

20

The reaction of the compound 3 with the compound 7 or 8 is carried out, for example, in an inert solvent or diluent, such as acetone, acetonitrile, dimethyl sulfoxide, dimethylformamide, N-methylpyrrolidone, etc., using a base (for example sodium

- 25 carbonate, potassium carbonate, sodium hydroxide, potassium hydroxide, sodium hydride, potassium hydride, etc.). The bases are generally used in equimolar amounts or in excess. Moreover, it may be advantageous to add a catalytic amount of a crown ether (for example 18-crown-6 or 15-crown-5). The reaction temperature
- 30 is generally in a range of from 0 to 80°C, preferably from 20 to 60°C.

The preparation of the compounds 7 and 8 is known to the person skilled in the art and described, for example, in Organikum, 17th

35 Edition, VEB Deutscher Verlag der Wissenschaften, Berlin 1988, p. 323ff, and Angewandte Chemie, 84 (1972) 295, and also Zh. Org. Khim 31 (1995) 601, Chem. Pharm. Bull. 36 (1988) 3134, Indian J. Chem., Sect. B, 31 (1992), 495.

- 40 The compound 4, obtained by reaction of 3 with 7, is subsequently oxamated using an O-alkenylhydroxylamine or a salt thereof. The preparation of the O-alkenylhydroxylamine is known to the person skilled in the art and described, for example, in Chem. Pharm. Bull. 91 (1983), 2601 and J. Am. Chem. Soc. 71 (1949), 3423.

45



## 20

Alternatively, it is possible to react a compound 3 directly with an oxime 8 to give a compound 1. All steps of route A, including the preparation of the intermediates or analogous intermediates, are described in EP 386 581A (for  $W = OCH_2$ ) and in EP 585 751A 5 (for  $W = C(R_{10})=NOCH_2$ ).

The starting material used for route B is likewise a compound 2. Reaction with the oxime 8 under the abovementioned etherification conditions gives a compound 6 ( $Y = NH_2$ ). The thiazoline ring is 10 then synthesized in the same manner as in route A.

Alternatively, a compound 2 is reacted with a compound 7 under the abovementioned etherification conditions to give a compound 5. This is oximated in the same manner as in route A, using 15 O-alkenylhydroxylamine or a salt thereof, giving the compound 6 into which, as already described, the thiazolone ring is then introduced.

Owing to the C=C or C=N double bonds, the compounds according to 20 the invention can be obtained in the preparation as E/Z isomer mixtures. These can be separated into the individual components in a customary manner, for example by crystallization or chromatography. Both the individual isomer compounds and their mixtures, and all enantiomers, racemates and diastereomers are 25 embraced by the invention.

The novel compounds 1 have excellent activity against a broad spectrum of phytopathogenic fungi, in particular from the classes of the Ascomycetes and Basidiomycetes, and they can be employed 30 as foliar- and soil-acting fungicides. Some of them have remarkably high systemic mobility and activity after soil application and in particular also after foliar application.

They are especially important for controlling a large number of 35 fungi on a variety of crop plants such as wheat, rye, barley, oats, rice, maize, grass, cotton, soybeans, coffee, sugar cane, grapevines, fruit species, ornamentals and vegetables such as cucumbers, beans, and cucurbits, and on the seeds of these plants.

40

Specifically, they are suitable for controlling the following plant diseases:

Erysiphe graminis (powdery mildew) in cereals,  
Erysiphe cichoracearum and Sphaerotheca fuliginea on cucurbits,  
45 Podosphaera leucotricha on apples,  
Uncinula necator on grapevines,

- Puccinia species on cereals,  
Rhizoctonia species on cotton and lawns,  
Ustilago species on cereals and sugar cane,  
Venturia inaequalis (scab) on apples,
- 5 Helminthosporium species on cereals,  
Septoria nodorum on wheat,  
Botrytis cinerea (gray mold) on strawberries, grapevines,  
Cercospora arachidicola on peanuts,  
Pseudocercospora herpotrichoides on wheat, barley,
- 10 Pyricularia oryzae on rice,  
Phytophthora infestans on potatoes and tomatoes,  
Fusarium and Verticillium species on a variety of plants,  
Plasmopara viticola on grapevines,  
Alternaria species on vegetables and fruit.
- 15
- The compounds 1 are applied by treating the fungi, or the plants, seeds, materials or the soil to be protected against fungal infection, with a fungicidally active amount of the active ingredients. Application is effected before or after infection of
- 20 the materials, plants or seeds by the fungi.

- They can be converted into the customary formulations, such as solutions, emulsions, suspensions, dusts, powders, pastes and granules. The use form depends on the specific intended use; in
- 25 any case, it should ensure fine and uniform distribution of the compounds 1 according to the invention. The formulations are prepared in a known manner, e.g. by extending the active ingredient with solvents and/or carriers, if desired using emulsifiers and dispersants, it also being possible to use other
- 30 organic solvents as auxiliary solvents if water is used as the diluent. Suitable auxiliaries for this purpose are essentially: solvents such as aromatics (e.g. xylene), chlorinated aromatics (e.g. chlorobenzenes), paraffins (e.g. mineral oil fractions), alcohols (e.g. methanol, butanol), ketones (e.g. cyclohexanone),
- 35 amines (e.g. ethanolamine, dimethylformamide) and water; carriers such as ground natural minerals (e.g. kaolins, clays, talc, chalk) and ground synthetic minerals (e.g. finely divided silica, silicates); emulsifiers such as nonionic and anionic emulsifiers (e.g. polyoxyethylene fatty alcohol ethers, alkylsulfonates and
- 40 arylsulfonates), and dispersants such as lignin-sulfite waste liquors and methylcellulose.

The fungicidal compositions generally comprise from 0.1 to 95, preferably from 0.5 to 90, % by weight of active ingredient.

Depending on the nature of the desired effect, the rates of application are from 0.01 to 3 kg of active ingredient per ha.

In the treatment of seed, amounts of active ingredient of from  
5 0.001 to 50 g, preferably 0.01 to 10 g, are generally required per kilogram of seed.

In the use form as fungicides, the compositions according to the invention can also exist together with other active ingredients,  
10 e.g. with herbicides, insecticides, growth regulators, fungicides, or else with fertilizers.

A mixture with fungicides frequently results in a widened fungicidal activity spectrum.

15

The compounds of the formula 1a are furthermore suitable for effectively controlling pests from the class of the insects, arachnids and nematodes. They can be employed as pesticides in crop protection and in the hygiene, stored-product and veterinary  
20 sector.

The harmful insects include, from the order of the lepidopterans (Lepidoptera), for example, *Agrotis ypsilon*, *Agrotis segetum*, *Alabama argillacea*, *Anticarsia gemmatalis*, *Argyresthia*  
25 *conjugella*, *Autographa gamma*, *Bupalus piniarius*, *Cacoecia murinana*, *Capua reticulana*, *Cheimatobia brumata*, *Choristoneura fumiferana*, *Choristoneura occidentalis*, *Cirphis unipuncta*, *Cydia pomonella*, *Dendrolimus pini*, *Diaphania nitidalis*, *Diatraea grandiosella*, *Earias insulana*, *Elasmopalpus lignosellus*,  
30 *Eupoecilia ambiguella*, *Evetria bouliana*, *Feltia subterranea*, *Galleria mellonella*, *Grapholitha funebrana*, *Grapholitha molesta*, *Heliothis armigera*, *Heliothis virescens*, *Heliothis zea*, *Hellula undalis*, *Hibernia defoliaria*, *Hyphantria cunea*, *Hyponomeuta malinellus*, *Keiferia lycopersicella*, *Lambdina fiscellaria*,  
35 *Laphygma exigua*, *Leucoptera coffeella*, *Leucoptera scitella*, *Lithocolletis blancardella*, *Lobesia botrana*, *Loxostege sticticalis*, *Lymantria dispar*, *Lymantria monacha*, *Lyonetia clerkella*, *Malacosoma neustria*, *Mamestra brassicae*, *Orgyia pseudotsugata*, *Ostrinia nubilalis*, *Panolis flammea*, *Pectinophora*  
40 *gossypiella*, *Peridroma saucia*, *Phalera bucephala*, *Phthorimaea operculella*, *Phyllocnistis citrella*, *Pieris brassicae*, *Plathypena scabra*, *Plutella xylostella*, *Pseudoplusia includens*, *Rhyacionia frustrana*, *Scrobipalpula absoluta*, *Sitotroga cerealella*, *Sparganothis pilleriana*, *Spodoptera frugiperda*, *Spodoptera*  
45 *littoralis*, *Spodoptera litura*, *Thaumatopoea pityocampa*, *Tortrix viridana*, *Trichoplusia ni*, *Zeiraphera canadensis*.

## 23

- From the order of the beetles (Coleoptera), for example, *Agrilus sinuatus*, *Agriotes lineatus*, *Agriotes obscurus*, *Amphimallus solstitialis*, *Anisandrus dispar*, *Anthonomus grandis*, *Anthonomus pomorum*, *Atomaria linearis*, *Blastophagus piniperda*, *Blitophaga*  
 5 *undata*, *Bruchus rufimanus*, *Bruchus pisorum*, *Bruchus lentis*, *Byctiscus betulae*, *Cassida nebulosa*, *Cerotoma trifurcata*, *Ceuthorrhynchus assimilis*, *Ceuthorrhynchus napi*, *Chaetocnema tibialis*, *Conoderus vespertinus*, *Crioceris asparagi*, *Diabrotica longicornis*, *Diabrotica 12-punctata*, *Diabrotica virgifera*,  
 10 *Epilachna varivestis*, *Epitrix hirtipennis*, *Eutinobothrus brasiliensis*, *Hylobius abietis*, *Hypera brunneipennis*, *Hypera postica*, *Ips typographus*, *Lema bilineata*, *Lema melanopus*, *Leptinotarsa decemlineata*, *Limonius californicus*, *Lissorhoptrus oryzophilus*, *Melanotus communis*, *Meligethes aeneus*, *Melolontha*  
 15 *hippocastani*, *Melolontha melolontha*, *Oulema oryzae*, *Otiorrhynchus sulcatus*, *Otiorrhynchus ovatus*, *Phaedon cochleariae*, *Phyllotreta chrysocephala*, *Phyllophaga* sp., *Phyllopertha horticola*, *Phyllotreta nemorum*, *Phyllotreta striolata*, *Popillia japonica*, *Sitona lineatus*, *Sitophilus granaria*.

## 20

- From the order of the dipterans (Diptera), for example, *Aedes aegypti*, *Aedes vexans*, *Anastrepha ludens*, *Anopheles maculipennis*, *Ceratitis capitata*, *Chrysomya bezziana*, *Chrysomya hominivorax*, *Chrysomya macellaria*, *Contarinia sorghicola*, *Cordylobia*  
 25 *anthropophaga*, *Culex pipiens*, *Dacus cucurbitae*, *Dacus oleae*, *Dasineura brassicae*, *Fannia canicularis*, *Gasterophilus intestinalis*, *Glossina morsitans*, *Haematobia irritans*, *Haplodiplosis equestris*, *Hylemyia platura*, *Hypoderma lineata*, *Liriomyza sativae*, *Liriomyza trifolii*, *Lucilia cuprina*, *Lucilia*  
 30 *sericata*, *Lycoria pectoralis*, *Mayetiola destructor*, *Musca domestica*, *Muscina stabulans*, *Oestrus ovis*, *Oscinella frit*, *Pegomya hyoscyami*, *Phorbia antiqua*, *Phorbia brassicae*, *Phorbia coarctata*, *Rhagoletis cerasi*, *Rhagoletis pomonella*, *Tabanus bovinus*, *Tipula oleracea*, *Tipula paludosa*.

## 35

From the order of the thrips (Thysanoptera), for example, *Frankliniella fusca*, *Frankliniella occidentalis*, *Frankliniella tritici*, *Scirtothrips citri*, *Thrips oryzae*, *Thrips palmi*, *Thrips tabaci*.

## 40

From the order of the hymenopterans (Hymenoptera), for example, *Athalia rosae*, *Atta cephalotes*, *Atta sexdens*, *Atta texana*, *Hoplocampa minuta*, *Hoplocampa testudinea*, *Monomorium pharaonis*, *Solenopsis geminata*, *Solenopsis invicta*.

## 45

From the order of the heteropterans (Heteroptera), for example, *Acrosternum hilare*, *Blissus leucopterus*, *Cyrtopeltis notatus*, *Dysdercus cingulatus*, *Dysdercus intermedius*, *Eurygaster integriceps*, *Euschistus impictiventris*, *Leptoglossus phyllopus*,  
 5 *Lygus lineolaris*, *Lygus pratensis*, *Nezara viridula*, *Piesma quadrata*, *Solubea insularis*, *Thyanta perditor*.

From the order of the homopterans (Homoptera), for example, *Acyrtosiphon onobrychis*, *Adelges laricis*, *Aphidula nasturtii*,  
 10 *Aphis fabae*, *Aphis pomi*, *Aphis sambuci*, *Bemisia tabaci*, *Brachycaudus cardui*, *Brevicoryne brassicae*, *Cerosipha gossypii*, *Dreyfusia nordmannianae*, *Dreyfusia piceae*, *Dysaphis radicola*, *Dysaulacorthum pseudosolani*, *Empoasca fabae*, *Macrosiphum avenae*, *Macrosiphum euphorbiae*, *Macrosiphon rosae*, *Megoura viciae*,  
 15 *Metopolophium dirhodum*, *Myzodes persicae*, *Myzus cerasi*, *Nephotettix cincticeps*, *Nilaparvata lugens*, *Pemphigus bursarius*, *Perkinsiella saccharicida*, *Phorodon humuli*, *Psylla mali*, *Psylla piri*, *Rhopalomyzus ascalonicus*, *Rhopalosiphum maidis*, *Sappahis mali*, *Schizaphis graminum*, *Schizoneura lanuginosa*, *Trialeurodes*  
 20 *vaporariorum*, *Viteus vitifolii*.

From the order of the termites (Isoptera), for example, *Calotermes flavicollis*, *Leucotermes flavipes*, *Reticulitermes lucifugus*, *Termes natalensis*.

25 From the order of the orthopterans (Orthoptera), for example, *Acheta domestica*, *Blatta orientalis*, *Blattella germanica*, *Forficula auricularia*, *Gryllotalpa gryllotalpa*, *Locusta migratoria*, *Melanoplus bivittatus*, *Melanoplus femur-rubrum*,  
 30 *Melanoplus mexicanus*, *Melanoplus sanguinipes*, *Melanoplus spretus*, *Nomadacris septemfasciata*, *Periplaneta americana*, *Schistocerca americana*, *Schistocerca peregrina*, *Stauronotus maroccanus*, *Tachycines asynamorus*.

35 From the class of the Arachnoidea, for example, arachnids (Acarina) such as *Amblyomma americanum*, *Amblyomma variegatum*, *Argas persicus*, *Boophilus annulatus*, *Boophilus decoloratus*, *Boophilus microplus*, *Brevipalpus phoenicis*, *Bryobia praetiosa*, *Dermacentor silvarum*, *Eotetranychus carpini*, *Eriophyes sheldoni*,  
 40 *Hyalomma truncatum*, *Ixodes ricinus*, *Ixodes rubicundus*, *Metatetranychus (Phanonychus) ulmi*, *Ornithodoros moubata*, *Otobius megnini*, *Paratetranychus pilosus*, *Dermanyssus gallinae*, *Phyllocoptiruta oleivora*, *Polyphagotarsonemus latus*, *Psoroptes ovis*, *Rhipicephalus appendiculatus*, *Rhipicephalus evertsi*,  
 45 *Sarcoptes scabiei*, *Tetranychus cinnabarinus*, *Tetranychus*

kanzawai, *Tetranychus pacificus*, *Tetranychus telarius*,  
*Tetranychus urticae*.

- From the class of the nematodes, for example, root knot  
 5 nematodes, e.g. *Meloidogyne hapla*, *Meloidogyne incognita*,  
*Meloidogyne javanica*, cyst-forming nematodes, e.g. *Globodera*  
*rostochiensis*, *Heterodera avenae*, *Heterodera glycines*, *Heterodera*  
*schachtii*, *Heterodera trifolii*, stem eelworms and foliar  
 nematodes, e.g. *Belonolaimus longicaudatus*, *Ditylenchus*  
 10  *destructor*, *Ditylenchus dipsaci*, *Helicotylenchus multicinctus*,  
*Longidorus elongatus*, *Radopholus similis*, *Rotylenchus robustus*,  
*Trichodorus primitivus*, *Tylenchorhynchus claytoni*,  
*Tylenchorhynchus dubius*, *Pratylenchus neglectus*, *Pratylenchus*  
*penetrans*, *Pratylenchus curvatus*, *Pratylenchus goodeyi*.

- 15 The active ingredients can be used as such, in the form of their  
 formulations or the use forms prepared therefrom, for example in  
 the form of ready-to-spray solutions, powders, suspensions or  
 dispersions, emulsions, oil dispersions, pastes, dusts, materials  
 20 for spreading or granules, by means of spraying, atomizing,  
 dusting, spreading or pouring. The use forms depend entirely on  
 the intended purposes; in any case, they should guarantee the  
 finest possible distribution of the active ingredients according  
 to the invention.

- 25 The concentrations of active ingredient in the ready-to-use  
 preparations can be varied within substantial ranges.

- In general, they are from 0.0001 to 10%, preferably from 0.01 to  
 30 1%.

- The active ingredients can also be used successfully in the  
 ultra-low-volume method (ULV), it being possible to apply  
 formulations with over 95% by weight of active ingredient, or  
 35 even the active ingredient without additives.

- Under field conditions, the rate of application of active  
 ingredient for controlling pests is 0.1 to 2.0, preferably 0.2 to  
 1.0 kg/ha.

- 40 Substances which are suitable for the preparation of  
 ready-to-spray solutions, emulsions, pastes or oil dispersions  
 are mineral oil fractions of medium to high boiling point such as  
 kerosene or diesel oil, furthermore coal-tar oils and oils of  
 45 vegetable or animal origin, aliphatic, cyclic and aromatic  
 hydrocarbons, e.g. benzene, toluene, xylene, paraffin,  
 tetrahydronaphthalene, alkylated naphthalenes or their

derivatives, methanol, ethanol, propanol, butanol, chloroform, carbon tetrachloride, cyclohexanol, cyclohexanone, chlorobenzene, isophorone, strongly polar solvents, e.g. dimethylformamide, dimethyl sulfoxide, N-methylpyrrolidone and water.

- 5 Aqueous use forms can be prepared from emulsion concentrates, pastes or wettable powders (sprayable powders, oil dispersions) by adding water. To prepare emulsions, pastes or oil dispersions, the substances, as such or dissolved in an oil or solvent, can be
- 10 homogenized in water by means of wetting agent, tackifier, dispersant or emulsifier. However, it is also possible to prepare concentrates composed of active ingredient, wetting agent, tackifier, dispersant or emulsifier and, if desired, solvent or oil, which are suitable for dilution with water.
- 15 Suitable surfactants are alkali metal, alkaline earth metal and ammonium salts of lignosulfonic acid, naphthalenesulfonic acid, phenolsulfonic acid, dibutyl naphthalenesulfonic acid, alkylarylsulfonates, alkyl sulfates, alkylsulfonates, fatty
- 20 alcohol sulfates and fatty acids and their alkali metal and alkaline earth metal salts, salts of sulfated fatty alcohol glycol ether, condensates of sulfonated naphthalene and naphthalene derivatives with formaldehyde, condensates of naphthalene or of naphthalenesulfonic acid with phenol and
- 25 formaldehyde, polyoxyethylene octylphenyl ether, ethoxylated isooctylphenol, octylphenol, nonylphenol, alkylphenyl polyglycol ethers, tributylphenyl polyglycol ether, alkylaryl polyether alcohols, isotridecyl alcohol, fatty alcohol/ethylene oxide condensates, ethoxylated castor oil, polyoxyethylene alkyl
- 30 ethers, ethoxylated polyoxypropylene, lauryl alcohol polyglycol ether acetal, sorbitol esters, lignin-sulfite waste liquors and methylcellulose.

- Powders, materials for spreading and dusts can be prepared by
- 35 mixing or concomitantly grinding the active substances with a solid carrier.

In general, the formulations comprise from 0.01 to 95% by weight, preferably from 0.1 to 90% by weight, of the active ingredient.

- 40 The active ingredients are employed in a purity of from 90% to 100%, preferably 95% to 100% (according to NMR spectrum).

Examples of the formulations are:

- 45 Granules, e.g. coated granules, impregnated granules and homogeneous granules; they can be prepared by binding the active ingredients onto solid carriers. Examples of solid carriers are

mineral earths such as silica gel, silicas, silicates, talc, kaolin, attaclay, limestone, lime, chalk, bole, loess, clay, dolomite, diatomaceous earth, calcium sulfate, magnesium sulfate, magnesium oxide, ground synthetic materials, fertilizers such as ammonium sulfate, ammonium phosphate, ammonium nitrate, ureas, and products of vegetable origin such as cereal meal, tree bark meal, wood meal and nut shell meal, cellulose powders, and other solid carriers.

10 Various types of oils, or herbicides, fungicides, other pesticides, or bactericides, can be added to the active ingredients, if appropriate also only just prior to use (tank mix). These agents can be admixed with the agents according to the invention in a weight ratio of 1:10 to 10:1.

15 In these use forms, the compositions according to the invention can also be present together with other active compounds, such as herbicides, insecticides, growth regulators and fungicides, or else be mixed and applied together with fertilizers. A mixture with fungicides frequently results in a widened fungicidal activity spectrum.

#### Example 1

25 Synthesis of 1,4-diacetylbenzene monooxime 9




35 25 g (0.154 mol) of 1,4-diacetylbenzene and 12.2 g (0.154 mol) of pyridine are dissolved in 80 ml of methanol. Under reflux, a solution of 10.7 g (0.154 mol) of hydroxylammonium chloride in 50 ml of water is added slowly. The mixture is stirred under  
40 reflux for 6 h and then at room temperature for 15 h. The mixture is then filtered, and the residue is washed three times with water and dried under reduced pressure. This gives 24.7 g (90%) of the product as colorless crystals. Melting point: 162 - 169°C.

45



## Example 2

## Synthesis of 10

Synthesis of 1,1-dimethyl-4-(*ortho*-tolyl)semicarbazide **6**:

At 10–15°C, 23.7 g (0.395 mol) of 1,1-dimethylhydrazine are slowly added to a solution of 50 g (0.376 mol) of *ortho*-tolyl isocyanate in 400 ml of absolute toluene. The mixture is stirred at room temperature for 16 h and then concentrated under reduced pressure, and the residue is suspended in 100 ml of cyclohexane. Filtration and washing of the residue with pentane gives 69.2 g (95%) of the product in the form of colorless crystals. M.p.: 134 – 136°C.

Synthesis of 5-chloro-2,4-dihydro-2-methyl-4-(*ortho*-tolyl)-3H-1,2,4-triazol-3-one **7**:

At 0°C, 106 g (0.359 mol) of triphosgene (bis(trichloromethyl) carbonate) are added to a solution of 69.2 g (0.358 mol) of 1,1-dimethyl-4-(*ortho*-tolyl)semicarbazide **6**. The reaction mixture is stirred at room temperature for 72 h and then concentrated under reduced pressure, and the residue is taken up in 150 ml of ethyl acetate. Cooling to 0°C results in 39.2 g of product precipitating out. The mother liquor is concentrated under

reduced pressure and taken up in 1.5 l of ethyl acetate. The mixture is washed three times with water, dried over sodium sulfate and concentrated under reduced pressure, giving a total of 49.5 g (65%) of product in the form of a colorless powder.

- 5 400 MHz-<sup>1</sup>H-NMR (CDCl<sub>3</sub>), δ [ppm]: 2.21 (s, 3H, Ar-CH<sub>3</sub>); 3.52 (s, 3H, N-CH<sub>3</sub>); 7.13-7.44 (m, 4H, Ar-H).

Synthesis of 2,4-dihydro-5-methoxy-2-methyl-4-(*ortho*-tolyl)-3H-1,2,4-triazol-3-one **8**:

10

79.2 g (0.44 mol) of a 30% strength sodium methoxide solution are added to a solution of 49.5 g (0.22 mol) of 5-chloro-2,4-dihydro-2-methyl-4-(*ortho*-tolyl)-3H-1,2,4-triazol-3-one **7** in 620 ml of a mixture of methanol and ethylene glycol dimethyl ether (1:1). The

- 15 mixture is stirred under reflux for 6 h and then concentrated under reduced pressure, and the residue is taken up in ethyl acetate. The mixture is washed three times with water, dried over sodium sulfate and concentrated under reduced pressure. The crude product is taken up in 100 ml of MTBE, admixed with 150 ml of  
20 hexane and cooled to 0°C. The precipitate is separated off and dried at 40°C under reduced pressure for 1 h. This gives 37.6 g (78%) of the product as a pale beige powder. M.p.: 127 - 130 °C.

Synthesis of 4-(*ortho*-bromomethylenepheryl)-2,4-dihydro-

- 25 5-methoxy-2-methyl-3H-1,2,4-triazol-3-one **10**:


A solution of 10.7 g (60 mmol) of *N*-bromosuccinimide in 150 ml of absolute 1,2-dichloroethane is added to a solution of 10.9 g (50 mmol) of 2,4-dihydro-5-methoxy-2-methyl-4-(*ortho*-tolyl)-

- 30 3H-1,2,4-triazol-3-one **8** and 0.4 g of azobisisobutyronitrile in 150 ml of absolute 1,2-dichloroethane. For 1 h, the mixture is stirred under reflux and irradiated with UV light. The mixture is then washed three times with water, dried over sodium sulfate and concentrated under reduced pressure. Purification of the residue  
35 by flash chromatography gives 4.1 g (28%) of the product as a colorless powder. M.p.: 113 - 115 °C.

Example 3: Synthesis of **11**


40

45



## 30


4.2 g (23 mmol) of 1,4-diacetylbenzene monooxime **9** are added to 0.62 g (26 mmol) of sodium hydride in 30 ml of absolute acetonitrile. The mixture is stirred under reflux for 1 h, and a solution of 7 g (23 mmol) of **10** in 50 ml of absolute acetonitrile **5** is then added at room temperature. The reaction mixture is stirred at room temperature for 15 h and then taken up in dilute NaCl solution and extracted three times with methyl *tert*-butyl ether. The combined organic phase is washed with water, dried over sodium sulfate and concentrated under reduced pressure. The **10** residue is purified by chromatography. This gives 7.2 g (79%) of the product as colorless crystals. Melting point 102 - 106°C.

Example 4: Synthesis of **12**

**25**

0.44 g (3.6 mmol) of O-2-trans-butenehydroxylammonium chloride and 1 g of molecular sieve 3 Å are added to a solution of 1.1 g (2.8 mmol) of **11** in 40 ml of methanol. The mixture is stirred at room temperature for 15 h. The molecular sieve is filtered off **30** and the reaction mixture is taken up in ethyl acetate, washed twice with water, dried with sodium sulfate and concentrated under reduced pressure. Chromatographic purification of the residue gives 840 ml (65%) of the residue as colorless crystals. Melting point 101 - 105°C.

**35**


Example 5: Synthesis of **13**

**45**

## 31

1.33 g (55 mmol) of sodium hydride are added to a solution of 8.3 g (50 mmol) of 2,5-dimethyl-4-hydroxyacetophenone in 120 ml of absolute DMF (dimethylformamide). The mixture is stirred at room temperature for 1 h, and a solution of 15 g (50 mmol) of 10  
5 in 80 ml of absolute DMF is then added, and the mixture is stirred at room temperature for 19 h. The mixture is then taken up in dilute NaCl solution and extracted three times with ethyl acetate. The combined organic phase is washed with water, dried over sodium sulfate and concentrated under reduced pressure.  
10 Chromatographic purification gives 13.9 g (73%) of the product.

## Example 6: Synthesis of 14



25 1 g (6 mmol) of 13 is dissolved in 40 ml of methanol. 0.42 g (2.9 mmol) of O-(3-chloroprop-2-ene)hydroxylammonium chloride and 1 g of molecular sieve 3 Å are added, and the mixture is stirred at room temperature for 15 h. The mixture is filtered and the  
30 filtrate is taken up in ethyl acetate, washed with water, dried with sodium sulfate and concentrated under reduced pressure, giving 1.1 g (89%) of the product (cf. Table No. 9 below).


The other compounds listed in Tables 2 and 3 below were obtained  
35 in a similar manner.

40

45

## 32

Table 2: Physical data of compounds of the formula:



No.	n	W	R <sub>2</sub>	R <sub>7</sub>	R <sub>8</sub>	R <sub>9</sub>	Analyt. data
15	1	-OCH <sub>2</sub> -	2-me-thyl	H	H	H	270MHz- <sup>1</sup> H-NMR (CDCl <sub>3</sub> ), δ (ppm): 2.22 (s, 3H); 2.24 (s, 3H); 3.41 (s, 3H); 3.91 (s, 3H); 4.67 (d, 2H); 5.10 (s, 2H); 5.20-5.40 (m, 3H); 5.98-6.18 (m, 1H); 6.78 (d, 1H); 7.22-7.64 (m, 6H).
20	2	-OCH <sub>2</sub> -	2-me-thyl	H	me-thyl	H	270MHz- <sup>1</sup> H-NMR (CDCl <sub>3</sub> ), δ (ppm): 1.74 (d, 3H); 2.19 (s, 3H); 2.21 (s, 3H); 3.42 (s, 3H); 3.92 (s, 3H); 4.60 (d, 2H); 5.10 (s, 2H); 5.71-5.86 (m, 2H); 6.78 (d, 1H); 7.19-7.63 (m, 6H).
25	3	-OCH <sub>2</sub> -	2-me-thyl	chloro	H	H	270MHz- <sup>1</sup> H-NMR (CDCl <sub>3</sub> ), δ (ppm): 2.20 (s, 3H); 2.22 (s, 3H); 3.41 (s, 3H); 3.93 (s, 3H); 4.71 (s, 2H); 5.19 (s, 2H); 5.18 (s, 1H); 5.22 (s, 1H); 6.78 (d, 1H); 7.21-7.64 (m, 6H).
30	4	-C(CH <sub>3</sub> )=NOCH <sub>2</sub> -	H	H	chloro	H	M.p. = 112-117°C
35	5	-C(CH <sub>3</sub> )=NOCH <sub>2</sub> -	H	H	H	H	270MHz- <sup>1</sup> H-NMR (CDCl <sub>3</sub> ), δ (ppm): 2.19 (s, 3H); 2.25 (s, 3H); 3.40 (s, 3H); 3.88 (s, 3H); 4.70 (d, 2H); 5.20-5.42 (m, 2H); 6.00-6.18 (m, 1H); 7.21-7.70 (m, 8H).
40	6	-C(CH <sub>3</sub> )=NOCH <sub>2</sub> -	H	H	methyl	H	M.p. = 101-105°C


## 33

5	7		-C(CH <sub>3</sub> )=NOCH <sub>2</sub> -	H	chloro	H	H	270MHz- <sup>1</sup> H-NMR (CDCl <sub>3</sub> ), δ (ppm): 2.19 (s, 3H); 2.24 (s, 3H); 3.40 (s, 3H); 3.94 (s, 3H); 4.76 (s, 2H); 5.23 (d, 2H); 5.40 (s, 1H); 5.45 (s, 1H); 7.21-7.61 (m, 8H).
10	8	1	-OCH <sub>2</sub> -	2-me- thyl	H	chloro	H	400MHz- <sup>1</sup> H-NMR (CDCl <sub>3</sub> ), δ (ppm): 2.19 (s, 3H); 2.23 (s, 3H); 3.42 (s, 3H); 3.92 (s, 3H); 4.62 (d, 2H); 5.10 (d, 2H); 6.15 (m, 1H); 6.26 (d, 1H); 6.79 (d, 1H); 7.22-7.63 (m, 7H).
15	9	2	-OCH <sub>2</sub> -	2,5-di- methyl	H	chloro	H	270MHz- <sup>1</sup> H-NMR (CDCl <sub>3</sub> ), δ (ppm): 2.14 (s, 3H); 2.17 (s, 3H); 2.24 (s, 3H); 3.42 (s, 3H); 3.93 (s, 3H); 4.61 (s, 2H); 5.02 (s, 2H); 6.03-6.11 (m, 1H); 6.60 (s, 1H); 6.98-7.64 (m, 7H).
20	10	2	-OCH <sub>2</sub> -	2,5-di- methyl	H	H	H	270MHz- <sup>1</sup> H-NMR (CDCl <sub>3</sub> ), δ (ppm): 2.17 (s, 6H); 2.26 (s, 3H); 3.44 (s, 3H); 3.90 (s, 3H); 4.62 (s, 2H); 5.06 (s, 2H); 5.97-6.13 (m, 1H); 6.60 (s, 1H); 6.99 (s, 1H); 7.20-7.65 (m, 6H).
25	11	2	-OCH <sub>2</sub> -	2,5-di- methyl	H	methyl	H	400MHz- <sup>1</sup> H-NMR (CDCl <sub>3</sub> ), δ (ppm): 1.72 (d, 3H); 2.15 (s, 3H); 2.20 (s, 3H); 2.30 (s, 3H); 3.44 (s, 3H); 3.97 (s, 3H); 4.95 (d, 2H); 5.07 (s, 2H); 5.60-5.83 (m, 2H); 6.60-7.61 (m, 6H).
30	12	2	-OCH <sub>2</sub> -	2,5-di- methyl	chloro	H	H	270MHz- <sup>1</sup> H-NMR (CDCl <sub>3</sub> ), δ (ppm): 2.07 (s, 3H); 2.15 (s, 3H); 2.26 (s, 3H); 3.42 (s, 3H); 3.95 (s, 3H); 4.85 (s, 2H); 5.05 (s, 2H); 5.14-5.23 (m, 2H); 6.60 (s, 1H); 7.21-7.63 (m, 5H).
35								
40								

34

5	13	1	-OCH <sub>2</sub> -	2-me- thyl	H	H	chloro	270MHz- <sup>1</sup> H-NMR (CDCl <sub>3</sub> ), δ (ppm): 2.18 (s, 3H); 2.23 (s, 3H); 3.41 (s, 3H); 3.90 (s, 3H); 4.93 (d, 2H); 5.09 (s, 2H); 6.02-6.21 (m, 2H); 6.88 (d, 1H); 7.21-7.64 (m, 7H).
10	14	2	-OCH <sub>2</sub> -	2,5-di- methyl	H	H	chloro	270MHz- <sup>1</sup> H-NMR (CDCl <sub>3</sub> ), δ (ppm): 2.07 (s, 3H); 2.18 (s, 3H); 2.25 (s, 3H); 3.43 (s, 3H); 3.92 (s, 3H); 4.82 (d, 2H); 6.00-7.64 (m, 8H).

Table 3: Physical data of compounds of formula:



No.	n	W	R2	A	Analyt. data	
30	15	1	-OCH <sub>2</sub> -	2-me- thyl	5-chlorothiophen- 2-yl	400MHz- <sup>1</sup> H-NMR (CDCl <sub>3</sub> ), δ (ppm): 2.19 (s, 3H); 2.23 (s, 3H); 3.42 (s, 3H); 3.89 (s, 3H); 5.19-5.30 (m, 4H); 6.79 (s, 1H); 6.84 (s, 1H); 7.23-7.70 (m, 8H).
	16	1	-OCH <sub>2</sub> -	2-me- thyl	2-(para-bromo- phenyl)oxazol- 4-yl	270MHz- <sup>1</sup> H-NMR (CDCl <sub>3</sub> ), δ (ppm): 1.59 (s, 3H); 2.23 (s, 3H), 3.41 (s, 3H), 3.95 (s, 3H); 5.10 (s, 2H); 5.21 (s, 2H); 6.80-7.95 (m, 12H).
40	17	1	-OCH <sub>2</sub> -	2-me- thyl	2-(para-chloro- phenyl)oxazol- 4-yl	270MHz- <sup>1</sup> H-NMR (CDCl <sub>3</sub> ), δ (ppm): 1.60 (s, 3H); 2.21 (s, 3H), 3.41 (s, 3H), 3.90 (s, 3H); 5.09 (s, 2H); 5.19 (s, 2H); 6.80 (d, 1H); 7.21-8.02 (m, 11H).

45

## 35

5	18	1	-OCH <sub>2</sub> -	2-me- thyl	thiophen-3-yl	270MHz- <sup>1</sup> H-NMR (CDCl <sub>3</sub> ), δ (ppm): 2.20 (s, 3H); 2.24 (s, 3H), 3.42 (s, 3H), 3.94 (s, 3H); 4.81 (d, 2H); 5.12 (s, 2H); 6.20-6.38 (m, 1H); 6.61-6.80 (m, 2H); 7.15-7.64 (m, 10H).
	19		-C(CH <sub>3</sub> )=NOCH <sub>2</sub> -	H	5-chlorothiophen- 2-yl	M.p. = 106-114°C
10	20		-C(CH <sub>3</sub> )=NOCH <sub>2</sub> -	H	2-(para-bromo- phenyl)oxazol- 4-yl	360MHz- <sup>1</sup> H-NMR (CDCl <sub>3</sub> ), δ (ppm): 2.17 (s, 3H); 2.21 (s, 3H), 3.35 (s, 3H), 3.43 (s, 3H); 4.80 (d, 2H); 5.21 (s, 2H); 7.18-7.95 (m, 13H).
	21		-C(CH <sub>3</sub> )=NOCH <sub>2</sub> -	H	thiophen-3-yl	M.p. = 109-126°C
15	22	2	-OCH <sub>2</sub> -	2,5- di- methyl	5-chlorothiophen- 2-yl	270MHz- <sup>1</sup> H-NMR (CDCl <sub>3</sub> ), δ (ppm): 2.17 (s, 3H); 2.20 (s, 3H); 2.30 (s, 3H); 3.41 (s, 3H); 3.96 (s, 3H); 5.02 (s, 2H); 5.19 (s, 2H); 6.80-7.66 (m, 8H).
	23	2	-OCH <sub>2</sub> -	2,5- di- methyl	2-(para-bromo- phenyl)oxazol- 4-yl	IR (cm <sup>-1</sup> ): 1725, 1616, 1501, 1479, 1414, 1402, 1389, 1325, 1247, 1149, 1073, 1037, 1010, 744, 733.
25	24	2	-OCH <sub>2</sub> -	2,5- di- methyl	thiophen-3-yl	270MHz- <sup>1</sup> H-NMR (CDCl <sub>3</sub> ), δ (ppm): 2.16 (s, 3H); 2.19 (s, 3H); 2.29 (s, 3H); 3.42 (s, 3H); 3.93 (s, 3H); 4.78 (d, 2H); 5.02 (s, 2H); 6.21-6.40 (m, 1H); 6.60-7.62 (m, 10H).
	25	2	-OCH <sub>2</sub> -	2,5- di- methyl	2-(para-chloro- phenyl)oxazol- 4-yl	M.p. = 92-117°C

## Use Examples

35

## Experiment 1 - Activity against mildew of wheat

Leaves of wheat seedlings c.v. "Frühgold" grown in pots were sprayed to runoff point with an aqueous preparation of active compound which had been prepared from a stock solution comprising 10% of active compound, 63% of cyclohexanone and 27% of emulsifier and, 24 hours after the spray coating had dried on, dusted with spores of mildew of wheat (*Erysiphe graminis forma specialis tritici*). The test plants were then placed in a greenhouse at 20-24°C and 60-90% relative atmospheric humidity.

45



After 7 days, the extent of the mildew development was determined visually in % infection of the total leaf area.

5	Active compound <sup>1)</sup>	% Infection of the leaves after application of an aqueous preparation comprising 63 ppm of active compound
	Active compound No. 1	5
	Active compound No. 2	10
10	Active compound No. 3	5
	Active compound No. 8	0
	Active compound No. 9	3
	Active compound No. 10	3
15	Active compound No. 11	0
	Active compound No. 12	5
	Active compound No. 13	10
	Active compound No. 14	5
	Active compound No. 15	15
20	Active compound No. 16	5
	Active compound No. 17	10
	Active compound No. 18	10
	Active compound No. 22	5
25	Active compound No. 23	3
	Active compound No. 24	5
	Active compound No. 25	5
30	Untreated	90

<sup>1)</sup> See Tables 2 and 3

#### Experiment 2 - Activity against *Plasmopara viticola*

- 35 Leaves of potted vines c.v. "Müller-Thurgau" were sprayed to runoff point with an aqueous preparation of active compound which had been prepared from a stock solution comprising 10% of active compound, 63% of cyclohexanone and 27% of emulsifier. The next day, the leaves were inoculated with an aqueous zoospore
- 40 suspension of *Plasmopara viticola*. The vines were then initially placed in a water-vapor-saturated chamber at 24°C for 48 hours and then in a greenhouse at 20-30°C for 5 days. After this time, the plants were once more placed in a humid chamber for 16 hours, to promote sporangiophore eruption. The extent of the development of
- 45 the infection on the underside of the leaves was then determined visually.

Active compound <sup>1)</sup>		% Infection of the leaves after application of an aqueous preparation comprising 63 ppm of active compound
5	Active compound No. 1	10
	Active compound No. 2	10
	Active compound No. 5	0
	Active compound No. 7	10
	Active compound No. 8	10
10	Active compound No. 9	1
	Active compound No. 10	0
	Active compound No. 11	0
	Active compound No. 12	0
	Active compound No. 13	15
15	Active compound No. 14	0
	Active compound No. 25	15
20	Untreated	85

<sup>1)</sup> See Tables 2 and 3

### Experiment 3 - Activity against *Pyricularia oryzae* (protective)

- 25 Leaves of rice seedlings c.v. "Tai-Nong 67" grown in pots were sprayed to runoff point with an aqueous preparation of active compound which had been prepared from a stock solution comprising 10% of active compound, 63% of cyclohexanone and 27% of emulsifier. The next day, the plants were inoculated with an aqueous sporangiophore suspension of *Pyricularia oryzae*. The test plants were then placed in climatized chambers at 22-24°C and 95-99% relative atmospheric humidity for 6 days. The extent of the development of the infection on the leaves was then determined visually.

35

Active compound <sup>1)</sup>		% Infection of the leaves after application of an aqueous preparation comprising 63 ppm of active compound
40	Active compound No. 4	15
	Active compound No. 5	5
	Active compound No. 6	10
	Active compound No. 7	5
	Active compound No. 8	5
45	Active compound No. 9	5
	Active compound No. 11	10
	Active compound No. 12	10

38

5	Active compound No. 13	5
	Active compound No. 14	5
	Active compound No. 15	10
	Active compound No. 22	5
	Active compound No. 23	5
10	Active compound No. 24	5
	Active compound No. 25	5
	Untreated	90

1) See Tables 2 and 3

15

20

25

30

35

40

45


ART 34 AMDT

We claim:

1. An oxime ether compound of the formula 1

5

10



where:

15

W is  $-\text{OCH}_2-$ ,  $-\text{C}(\text{R}_{10})=\text{N}-\text{O}-\text{CH}_2-$ ;

X is halogen,  $\text{C}_1$ - $\text{C}_4$ -alkyl,  $\text{C}_1$ - $\text{C}_4$ -alkoxy;

20

$\text{R}_1$  is H,  $\text{C}_1$ - $\text{C}_4$ -alkyl, halogen, nitro, CN, halo- $\text{C}_1$ - $\text{C}_4$ -alkyl,  $\text{C}_1$ - $\text{C}_4$ -alkoxy;

$\text{R}_2$  is H,  $\text{C}_1$ - $\text{C}_4$ -alkyl, halogen, nitro, CN, halo- $\text{C}_1$ - $\text{C}_4$ -alkyl,  $\text{C}_1$ - $\text{C}_4$ -alkoxy;

25

n is 1 or 2;

$\text{R}_3$  is H,  $\text{C}_1$ - $\text{C}_4$ -alkyl;

30

$\text{R}_5$  is H,  $\text{C}_1$ - $\text{C}_4$ -alkyl,  $\text{C}_2$ - $\text{C}_4$ -alkenyl;

$\text{R}_6$  is H,  $\text{C}_1$ - $\text{C}_4$ -alkyl,  $\text{C}_1$ - $\text{C}_4$ -haloalkyl,  $\text{C}_2$ - $\text{C}_4$ -alkenyl, aryl;

35

$\text{R}_7$  is H, halogen,  $\text{C}_1$ - $\text{C}_6$ -alkyl,  $\text{C}_1$ - $\text{C}_6$ -haloalkyl,  $\text{C}_2$ - $\text{C}_6$ -alkenyl,  $\text{C}_2$ - $\text{C}_6$ -haloalkenyl,  $\text{C}_3$ - $\text{C}_6$ -cycloalkyl,  $\text{C}_3$ - $\text{C}_6$ -halocycloalkyl, unsubstituted or substituted aryl;

40

$\text{R}_8$  is H, halogen,  $\text{C}_1$ - $\text{C}_6$ -alkyl,  $\text{C}_1$ - $\text{C}_6$ -haloalkyl,  $\text{C}_2$ - $\text{C}_6$ -alkenyl,  $\text{C}_2$ - $\text{C}_6$ -haloalkenyl,  $\text{C}_3$ - $\text{C}_6$ -cycloalkyl,  $\text{C}_3$ - $\text{C}_6$ -halocycloalkyl, unsubstituted or substituted aryl, or

45

- 5  $R_7$  and  $R_8$ , together with the carbon atoms to which they are attached, form an unsaturated heterocycle having 5 or 6 ring atoms and one or two heteroatoms, independently of one another selected from nitrogen, oxygen and sulfur, which heterocycle may be substituted by one or two radicals which, independently of one another, are selected from the group consisting of  $C_1$ - $C_4$ -alkyl, halogen, nitro, CN, halo- $C_1$ - $C_4$ -alkyl, OH,  $C_1$ - $C_4$ -alkoxy, unsubstituted or substituted aryl,  $C_2$ - $C_4$ -alkenyl,
- 10 halo- $C_2$ - $C_4$ -alkenyl,  $C_2$ - $C_4$ -alkynyl, halo- $C_2$ - $C_4$ -alkynyl;
- $R_9$  is H, halogen,  $C_1$ - $C_6$ -alkyl,  $C_1$ - $C_6$ -haloalkyl,  $C_2$ - $C_6$ -alkenyl,  $C_2$ - $C_6$ -haloalkenyl,  $C_3$ - $C_6$ -cycloalkyl,  $C_3$ - $C_6$ -halocycloalkyl, unsubstituted or substituted aryl;
- 15  $R_{10}$  is H, halogen,  $C_1$ - $C_4$ -alkyl.
2. A compound of the formula 1 as claimed in claim 1 where:
- 20 W is  $-OCH_2-$ ,  $-C(R_{10})=N-O-CH_2$ ;
- X is halogen,  $C_1$ - $C_4$ -alkyl,  $C_1$ - $C_4$ -alkoxy;
- 25  $R_1$  is H,  $C_1$ - $C_4$ -alkyl, halogen, halo- $C_1$ - $C_4$ -alkyl;
- $R_2$  is H,  $C_1$ - $C_4$ -alkyl, halogen, halo- $C_1$ - $C_4$ -alkyl;
- $R_3$  is H,  $C_1$ - $C_4$ -alkyl;
- 30 n is 1 or 2;
- $R_5$  is H or  $C_1$ - $C_4$ -alkyl;
- 35  $R_6$  is H,  $C_1$ - $C_4$ -alkyl, halo- $C_1$ - $C_4$ -alkyl;
- $R_7$  is H, halogen,  $C_1$ - $C_6$ -alkyl, halo- $C_1$ - $C_6$ -alkyl,  $C_3$ - $C_6$ -cycloalkyl,  $C_3$ - $C_6$ -halocycloalkyl, phenyl;
- 40  $R_8$  is H, halogen,  $C_1$ - $C_6$ -alkyl, halo- $C_1$ - $C_6$ -alkyl,  $C_3$ - $C_6$ -cycloalkyl,  $C_3$ - $C_6$ -halocycloalkyl,  $C_2$ - $C_6$ -alkenyl, phenyl which may be substituted by one or two halogen or  $C_1$ - $C_4$ -alkyl; or

- 45  $R_7$  and  $R_8$ , together with the carbon atoms to which they are attached, form an unsaturated heterocycle having 5 or 6 ring atoms and one or two heteroatoms, independently of one another selected from nitrogen, oxygen and sulfur,

## Translation of amended sheet annexed to IPER

- 41 -  
- 3 -

which heterocycle may be substituted by one or two radicals which, independently of one another, are C<sub>1</sub>-C<sub>4</sub>-alkyl, halogen, halo-C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy and phenyl which may be substituted by one or two halogen or C<sub>1</sub>-C<sub>4</sub>-alkyl;

5

R<sub>9</sub> is H, halogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, halo-C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, C<sub>3</sub>-C<sub>6</sub>-halocycloalkyl, phenyl;

10

R<sub>10</sub> is H, halogen, C<sub>1</sub>-C<sub>4</sub>-alkyl.

3. A compound of the formula 1 as claimed in claim 1 or 2 where:

15

W is -OCH<sub>2</sub>-, -C(R<sub>10</sub>)=N-O-CH<sub>2</sub>;

X is halogen, C<sub>1</sub>-C<sub>4</sub>-alkoxy;

R<sub>1</sub> is H, C<sub>1</sub>-C<sub>4</sub>-alkyl;

20

R<sub>2</sub> is H, C<sub>1</sub>-C<sub>4</sub>-alkyl;

n is 1 or 2;

R<sub>3</sub> is C<sub>1</sub>-C<sub>4</sub>-alkyl;

25

R<sub>5</sub> is H, C<sub>1</sub>-C<sub>4</sub>-alkyl;

R<sub>6</sub> is H, C<sub>1</sub>-C<sub>4</sub>-alkyl;

30

R<sub>7</sub> is H, halogen, C<sub>1</sub>-C<sub>6</sub>-alkyl;

R<sub>8</sub> is H, halogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>2</sub>-C<sub>6</sub>-alkenyl; or

35

R<sub>7</sub> and R<sub>8</sub>, together with the carbon atoms to which they are attached, form a thiophenyl, furanyl, oxazolyl or thiazolyl radical, where these groups may have one or two substituents which are selected independently of one another from the group consisting of C<sub>1</sub>-C<sub>4</sub>-alkyl, halogen and phenyl which may be substituted by one or two

40

halogen;

R<sub>9</sub> is H, halogen, C<sub>1</sub>-C<sub>6</sub>-alkyl;

R<sub>10</sub> is H, C<sub>1</sub>-C<sub>4</sub>-alkyl.

45

4. A compound of the formula 1 as claimed in any of the preceding claims where:

- 5       W    is  $-\text{OCH}_2-$ ,  $-\text{C}(\text{R}_{10})=\text{N}-\text{O}-\text{CH}_2$ ;
- X    is  $\text{C}_1$ - $\text{C}_4$ -alkoxy;
- $\text{R}_1$    is H;
- 10        $\text{R}_2$    is H,  $\text{C}_1$ - $\text{C}_4$ -alkyl;
- n    is 1 or 2;
- 15        $\text{R}_3$    is  $\text{C}_1$ - $\text{C}_4$ -alkyl;
- $\text{R}_5$    is H,  $\text{C}_1$ - $\text{C}_4$ -alkyl;
- $\text{R}_6$    is H,  $\text{C}_1$ - $\text{C}_4$ -alkyl;
- 20        $\text{R}_7$    is H, halogen;
- $\text{R}_8$    is H,  $\text{C}_1$ - $\text{C}_4$ -alkyl, halogen; or
- 25        $\text{R}_7$  and  $\text{R}_8$ , together with the carbon atoms to which they are attached, form a thiophenyl or oxazolyl radical, where these groups may be substituted by one or two halogen or phenyl and the phenyl may be substituted by one or two halogen;
- 30        $\text{R}_9$    is H, halogen;
- $\text{R}_{10}$    is H,  $\text{C}_1$ - $\text{C}_4$ -alkyl.
5. A compound of the formula 1 where:
- 35       W    is  $-\text{OCH}_2-$ ,  $-\text{C}(\text{R}_{10})=\text{N}-\text{O}-\text{CH}_2-$ ;
- X    is halogen,  $\text{C}_1$ - $\text{C}_4$ -alkyl,  $\text{C}_1$ - $\text{C}_4$ -alkoxy;
- 40        $\text{R}_1$    is H,  $\text{C}_1$ - $\text{C}_4$ -alkyl, halogen, nitro, CN, halo- $\text{C}_1$ - $\text{C}_4$ -alkyl,  $\text{C}_1$ - $\text{C}_4$ -alkoxy;
- $\text{R}_2$    is H,  $\text{C}_1$ - $\text{C}_4$ -alkyl, halogen, nitro, CN, halo- $\text{C}_1$ - $\text{C}_4$ -alkyl,  $\text{C}_1$ - $\text{C}_4$ -alkoxy;
- 45       n    is 1 or 2;


- R<sub>3</sub> is H, C<sub>1</sub>-C<sub>4</sub>-alkyl;
- R<sub>5</sub> is H, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>2</sub>-C<sub>4</sub>-alkenyl;
- 5 R<sub>6</sub> is H, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkyl, C<sub>2</sub>-C<sub>4</sub>-alkenyl, aryl;
- R<sub>7</sub> is H, halogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkyl,  
C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>2</sub>-C<sub>6</sub>-haloalkenyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl,  
C<sub>3</sub>-C<sub>6</sub>-halocycloalkyl, unsubstituted or substituted aryl;
- 10 R<sub>8</sub> is H, halogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkyl,  
C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>2</sub>-C<sub>6</sub>-haloalkenyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl,  
C<sub>3</sub>-C<sub>6</sub>-halocycloalkyl, unsubstituted or substituted aryl;
- 15 R<sub>9</sub> is H, halogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkyl,  
C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>2</sub>-C<sub>6</sub>-haloalkenyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl,  
C<sub>3</sub>-C<sub>6</sub>-halocycloalkyl, unsubstituted or substituted aryl;
- R<sub>10</sub> is H, halogen, C<sub>1</sub>-C<sub>4</sub>-alkyl.
- 20 6. The use of the compounds of the formula 1 as claimed in any  
of claims 1 to 5 as fungicides or for controlling pests.
- 25 7. A fungicide, comprising solid and/or liquid carriers and a  
fungicidally effective amount of at least one compound of the  
formula 1 as claimed in claim 1.
- 30 8. A method for controlling fungi, wherein the fungi or the  
materials, plants, seeds or the soil threatened by fungal  
attack is treated with a fungicidally effective amount of at  
least one compound of the formula 1 as claimed in claim 1.
- 35 9. A composition for controlling pests, comprising inert  
additives and a pesticidally effective amount of at least one  
compound of the formula 1 as claimed in claim 1.
- 40 10. A method for controlling pests, wherein the pests and/or  
their habitat are treated with a pesticidally effective  
amount of at least one compound of the formula 1 as claimed  
in claim 1.



44

6

11. A compound of the formula 6



in which

15 W, R<sub>1</sub>, R<sub>2</sub>, R<sub>5</sub>, R<sub>6</sub>, R<sub>7</sub>, R<sub>8</sub>, R<sub>9</sub> and n are as defined in any of claims 1 to 5 and Y is NH<sub>2</sub>.

20

25

30


35

40

45

## Abstract

Unsaturated oxime ether compounds of the formula 1



where:

- 15 W is  $-\text{OCH}_2-$ ,  $-\text{C}(\text{R}_{10})=\text{N}-\text{O}-\text{CH}_2-$ ;
- X is halogen,  $\text{C}_1$ - $\text{C}_4$ -alkyl,  $\text{C}_1$ - $\text{C}_4$ -alkoxy;
- 20  $\text{R}_1$  is H,  $\text{C}_1$ - $\text{C}_4$ -alkyl, halogen, nitro, CN, halo- $\text{C}_1$ - $\text{C}_4$ -alkyl,  $\text{C}_1$ - $\text{C}_4$ -alkoxy;
- $\text{R}_2$  is H,  $\text{C}_1$ - $\text{C}_4$ -alkyl, halogen, nitro, CN, halo- $\text{C}_1$ - $\text{C}_4$ -alkyl,  $\text{C}_1$ - $\text{C}_4$ -alkoxy;
- 25 n is 1 or 2;
- $\text{R}_3$  is H,  $\text{C}_1$ - $\text{C}_4$ -alkyl;
- 30  $\text{R}_5$  is H,  $\text{C}_1$ - $\text{C}_4$ -alkyl,  $\text{C}_2$ - $\text{C}_4$ -alkenyl;
- $\text{R}_6$  is H,  $\text{C}_1$ - $\text{C}_4$ -alkyl,  $\text{C}_1$ - $\text{C}_4$ -haloalkyl,  $\text{C}_2$ - $\text{C}_4$ -alkenyl, aryl;
- 35  $\text{R}_7$  is H, halogen,  $\text{C}_1$ - $\text{C}_6$ -alkyl,  $\text{C}_1$ - $\text{C}_6$ -haloalkyl,  $\text{C}_2$ - $\text{C}_6$ -alkenyl,  $\text{C}_2$ - $\text{C}_6$ -haloalkenyl,  $\text{C}_3$ - $\text{C}_6$ -cycloalkyl,  $\text{C}_3$ - $\text{C}_6$ -halocycloalkyl, unsubstituted or substituted aryl;
- $\text{R}_8$  is H, halogen,  $\text{C}_1$ - $\text{C}_6$ -alkyl,  $\text{C}_1$ - $\text{C}_6$ -haloalkyl,  $\text{C}_2$ - $\text{C}_6$ -alkenyl,  $\text{C}_2$ - $\text{C}_6$ -haloalkenyl,  $\text{C}_3$ - $\text{C}_6$ -cycloalkyl,  $\text{C}_3$ - $\text{C}_6$ -halocycloalkyl, unsubstituted or substituted aryl, or
- 40  $\text{R}_7$  and  $\text{R}_8$ , together with the carbon atoms to which they are attached, form an unsaturated heterocycle having 5 or 6 ring atoms and one or two heteroatoms, independently of one another selected from nitrogen, oxygen and sulfur, which heterocycle may be substituted by one or two radicals which, independently of one another, are selected from the group
- 45

~~46~~  
45

consisting of C<sub>1</sub>-C<sub>4</sub>-alkyl, halogen, nitro, CN, halo-C<sub>1</sub>-C<sub>4</sub>-alkyl, OH, C<sub>1</sub>-C<sub>4</sub>-alkoxy, unsubstituted or substituted aryl, C<sub>2</sub>-C<sub>4</sub>-alkenyl, halo-C<sub>2</sub>-C<sub>4</sub>-alkenyl, C<sub>2</sub>-C<sub>4</sub>-alkynyl, halo-C<sub>2</sub>-C<sub>4</sub>-alkynyl;

5

R<sub>9</sub> is H, halogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>2</sub>-C<sub>6</sub>-haloalkenyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl, C<sub>3</sub>-C<sub>6</sub>-halocycloalkyl, unsubstituted or substituted aryl;

10 R<sub>10</sub> is H, halogen, C<sub>1</sub>-C<sub>4</sub>-alkyl,

are described. The compounds according to the invention are suitable for controlling harmful fungi and animal pests.

15

20

25

30

35

40

45

# Declaration, Power of Attorney



26474

PATENT & TRADEMARK OFFICE

Page 1 of 6

0050/050730

We (I), the undersigned inventor(s), hereby declare(s) that:

My residence, post office address and citizenship are as stated below next to my name,

We (I) believe that we are (I am) the original, first, and joint (sole) inventor(s) of the subject matter which is claimed and for which a patent is sought on the invention entitled

Unsaturated oxime ethers and the use thereof for control of harmful fungi and veterinary pests

the specification of which

☒ is attached hereto.

☐ was filed on \_\_\_\_\_ as

Application Serial No. \_\_\_\_\_

and amended on \_\_\_\_\_.

☒ was filed as PCT international application

Number PCT/EP00/09000

on September 14, 2000

and was amended under PCT Article 19

on \_\_\_\_\_ (if applicable).

We (I) hereby state that we (I) have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

We (I) acknowledge the duty to disclose information known to be material to the patentability of this application as defined in Section 1.56 of Title 37 Code of Federal Regulations.

We (I) hereby claim foreign priority benefits under 35 U.S.C. § 119(a)-(d) or § 365(b) of any foreign application(s) for patent or inventor's certificate, or § 365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or PCT International application having a filing date before that of the application on which priority is claimed. Prior Foreign Application(s)

Application No.	Country	Day/Month/Year	Priority Claimed
19944258.4	Germany	15 September 1999	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No

pduskpet001 - 25

We (I) hereby claim the benefit under Title 35, United States Codes, § 119(e) of any United States provisional application(s) listed below.

\_\_\_\_\_  
(Application Number)

\_\_\_\_\_  
(Filing Date)

\_\_\_\_\_  
(Application Number)

\_\_\_\_\_  
(Filing Date)

We (I) hereby claim the benefit under 35 U.S.C. § 120 of any United States application(s), or § 365(c) of any PCT International application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of 35 U.S.C. § 112, I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR § 1.56 which became available between the filing date of the prior application and the national or PCT International filing date of this application.

Application Serial No.	Filing Date	Status (pending, patented, abandoned)
_____	_____	_____
_____	_____	_____
_____	_____	_____
_____	_____	_____

And we (I) hereby appoint Messrs. **HERBERT. B. KEIL**, Registration Number 18,967; and **RUSSEL E. WEINKAUF**, Registration Number 18,495; the address of both being Messrs. Keil & Weinkauf, 1101 Connecticut Ave., N.W., Washington, D.C. 20036 (telephone 202-659-0100), our attorneys, with full power of substitution and revocation, to prosecute this application, to make alterations and amendments therein, to sign the drawings, to receive the patent, and to transact all business in the Patent Office connected therewith.

We (I) declare that all statements made herein of our (my) own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

1 - 00 Markus Gewehr  
NAME OF INVENTOR

Goethestraße 21  
56288 Kastellaun DEX  
Germany  
Citizen of: Germany  
Post Office Address: same as residence

Markus Gewehr  
Signature of Inventor

Date January 18, 2002

2 - 00 Hubert Sauter  
NAME OF INVENTOR

Neckarpromenade 20  
68167 Mannheim DEX  
Germany  
Citizen of: Germany  
Post Office Address: same as residence

Hubert Sauter  
Signature of Inventor

Date January 18, 2002

3 - 00 Bernd Müller  
NAME OF INVENTOR

Jean-Ganss-Str.21  
67227 Frankenthal DEX  
Germany  
Citizen of: Germany  
Post Office Address: same as residence

Bernd Müller  
Signature of Inventor

Date January 18, 2002

4 - 00 Andreas Gypser  
NAME OF INVENTOR

B 4.4  
68159 Mannheim DEX  
Germany  
Citizen of: Germany  
Post Office Address: same as residence

Andreas Gypser  
Signature of Inventor

Date January 18, 2002

5-00 Wassilios Grammenos  
NAME OF INVENTOR

Wassilios Grammenos  
Signature of Inventor

Date January 18, 2002

Borsigstr.5  
67063 Ludwigshafen DEX  
Germany  
Citizen of: Greece  
Post Office Address: same as residence

6-00 Jordi Tormo i Blasco  
NAME OF INVENTOR

Jordi Tormo i Blasco  
Signature of Inventor

Date January 18, 2002

Mühlweg 47  
67117 Limburgerhof DEX  
Germany  
Citizen of: Spain  
Post Office Address: same as residence

7-00 Arne Ptöck  
NAME OF INVENTOR

Arne Ptöck  
Signature of Inventor

Date January 18, 2002

Tiroler Str. 33  
67065 Ludwigshafen DEX  
Germany  
Citizen of: Germany  
Post Office Address: same as residence

8-00 Oliver Cullmann  
NAME OF INVENTOR

Oliver Cullmann  
Signature of Inventor

Date January 18, 2002

Heinrich-Heine-Str.27  
68199 Mannheim DEX  
Germany  
Citizen of: Germany  
Post Office Address: same as residence

9 - 00 Thomas Grote  
NAME OF INVENTOR

Thomas Grote  
Signature of Inventor

Date January 18, 2002

Im Hoehnhausen  
67157 Wachenheim DEX  
Germany  
Citizen of: Germany  
Post Office Address: same as residence

10 - 00 Eberhard Ammermann  
NAME OF INVENTOR

Eberhard Ammermann  
Signature of Inventor

Date January 18, 2002

Von-Gagern-Str.2  
64646 Heppenheim DEX  
Germany  
Citizen of: Germany  
Post Office Address: same as residence

11 - 00 Siegfried Strathmann  
NAME OF INVENTOR

Siegfried Strathmann  
Signature of Inventor

Date January 18, 2002

Donnersbergstr.9  
67117 Limburgerhof DEX  
Germany  
Citizen of: Germany  
Post Office Address: same as residence

12 - 00 Gisela Lorenz  
NAME OF INVENTOR

Gisela Lorenz  
Signature of Inventor

Date January 18, 2002

Erlenweg 13  
67434 Neustadt DEX  
Germany  
Citizen of: Germany  
Post Office Address: same as residence



**Declaration**

Page 6 of 6

0050/050730

13-00 Volker Harries  
NAME OF INVENTOR

Volker Harries

Signature of Inventor

Date January 18, 2002

Immengärtenweg 29e

67227 Frankenthal D E X

Germany

Citizen of: Germany

Post Office Address: same as residence